



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01H 5/00, 5/10, A23D 7/00, A23K 1/14, C07H 21/04, C07K 4/10, C12N 15/04, 15/63, 15/82	A1	(11) International Publication Number: WO 99/02030 (43) International Publication Date: 21 January 1999 (21.01.99)
(21) International Application Number: PCT/US98/14528 (22) International Filing Date: 10 July 1998 (10.07.98) (30) Priority Data: 60/052,249 11 July 1997 (11.07.97) US (71) Applicant (for all designated States except US): WASHINGTON STATE UNIVERSITY RESEARCH FOUNDATION [US/US]; N.E. 1615 Eastgate Boulevard, Pullman, WA 99164-1802 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BOHLMANN, Joerg [DE/DE]; Sophienstrasse 21, D-0745 Jena (DE). STEELE, Christopher, L. [US/US]; 617 Campbell Street, Ardmore, OK 973401 (US). CROTEAU, Rodney, B. [US/US]; N.E. 1835 Valley Road, Pullman, WA 99163 (US). (74) Agent: SHELTON, Dennis, K.; Christensen O'Connor Johnson & Kindness PLLC, Suite 2800, 1420 Fifth Avenue, Seattle, WA 98101 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: MONOTERPENE SYNTHASES FROM GRAND FIR (<i>ABIES GRANDIS</i>)		
(57) Abstract <p>cDNAs encoding myrcene synthase, (–)-limonene synthase and (–)-pinene synthase from Grand fir (<i>Abies grandis</i>) haven been isolated and sequenced, and the corresponding amino acid sequences have been determined. Accordingly, isolated DNA sequences (SEQ ID NO:1; SEQ ID NO:3 and SEQ ID NO:5) are provided which code for the expression of myrcene synthase (SEQ ID NO:2), (–)-pinene synthase (SEQ ID NO:4) and (–)-limonene synthase (SEQ ID NO:6) , respectively, from Grand fir (<i>Abies grandis</i>). In other aspects, replicable recombinant cloning vehicles are provided which code for myrcene synthase, (–)-limonene synthase and (–)-pinene synthase, or for a base sequence sufficiently complementary to at least a portion of myrcene synthase, (–)-limonene synthase or (–)-pinene synthase DNA or RNA to enable hybridization therewith. In yet other aspects, modified host cells are provided that have been transformed, transfected, infected and/or injected with a recombinant cloning vehicle and/or DNA sequence encoding myrcene synthase, (–)-limonene synthase or (–)-pinene synthase. Thus, systems and methods are provided for the recombinant expression of the aforementioned recombinant monoterpene synthases that may be used to facilitate their production, isolation and purification in significant amounts. Recombinant myrcene synthase, (–)-limonene synthase or (–)-pinene synthase may be used to obtain expression or enhanced expression of myrcene synthase, (–)-limonene synthase or (–)-pinene synthase in plants in order to enhance the production of monoterpenoids, or may be otherwise employed for the regulation or expression of myrcene synthase, (–)-limonene synthase and (–)-pinene synthase, or the production of their products.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MONOTERPENE SYNTHASES FROM GRAND FIR (*Abies grandis*)

Field of the Invention

5 The present invention relates to nucleic acid sequences which code for monoterpene synthases from gymnosperm plant species, in particular from Grand fir (*Abies grandis*), including limonene synthase, myrcene synthase, and pinene synthase, to vectors containing the sequences, to host cells containing the sequences, to plant seeds expressing the sequences and to methods of producing recombinant monoterpene synthases and their mutants.

Background of the Invention

10 Chemical defense of conifer trees against bark beetles and their associated fungal pathogens relies primarily upon constitutive and inducible oleoresin biosynthesis (Johnson, M.A., and Croteau, R. (1987) in *Ecology and Metabolism of Plant Lipids* (Fuller, G., and Nes, W.D., eds.) pp. 76-91, American Chemical Society Symposium Series 325, Washington, DC; Gijzen, M., Lewinsohn, E., Savage, T.J., and Croteau, R.B. (1993) in *Bioactive Volatile Compounds from Plants* (Teranishi, R., Buttery, R.G., and Sugisawa, H., eds.) pp. 8-22, American Chemical Society Symposium Series 525, Washington, DC). This defensive secretion is a complex mixture of monoterpene and sesquiterpene olefins (turpentine) and diterpene resin acids (rosin) that is synthesized constitutively in the epithelial cells of specialized structures, such as resin ducts and blisters or, in the case of induced oleoresin formation, in undifferentiated cells surrounding wound sites (Lewinsohn, E., Gijzen, M., Savage, T.J., and Croteau, R. (1991) *Plant Physiol.* **96**:38-43). The volatile fraction of conifer oleoresin, which is toxic to both bark beetles and their fungal

associates (Raffa, K.F., Berryman, A.A., Simasko, J., Teal, W., and Wong, B.L. (1985) *Environ. Entomol.* **14**:552-556), may consist of up to 30 different monoterpenes (Lewinsohn, E., Savage, T.J., Gijzen, M., and Croteau, R. (1993) *Phytochem. Anal.* **4**:220-225), including acyclic types (e.g., myrcene), monocyclic types (e.g., limonene) and bicyclic types (e.g., pinenes) (FIGURE 1). Although the oleoresin is toxic, many bark beetle species nevertheless employ turpentine volatiles in host selection and can convert various monoterpene components into aggregation or sex pheromones to promote coordinated mass attack of the host (Gijzen, M., Lewinsohn, E., Savage, T.J., and Croteau, R.B. (1993) in *Bioactive Volatile Compounds from Plants* (Teranishi, R., Buttery, R.G., and Sugisawa, H., eds.) pp. 8-22, American Chemical Society Symposium Series 525, Washington, DC; Byers, J.A. (1995) in *Chemical Ecology of Insects 2* (Cardé, R.T., and Bell, W.J., eds.) pp. 154-213, Chapman and Hall, New York). In Grand fir (*Abies grandis*), increased formation of oleoresin monoterpenes, sesquiterpenes and diterpenes is induced by bark beetle attack (Lewinsohn, E., Gijzen, M., Savage, T.J., and Croteau, R. (1991) *Plant Physiol.* **96**:38-43; Raffa, K.F., and Berryman, A.A. (1982) *Can. Entomol.* **114**:797-810; Lewinsohn, E., Gijzen, M., and Croteau, R. (1991) *Plant Physiol.* **96**:44-49), and this inducible defense response is mimicked by mechanically wounding sapling stems (Lewinsohn, E., Gijzen, M., Savage, T.J., and Croteau, R. (1991) *Plant Physiol.* **96**:38-43; Lewinsohn, E., Gijzen, M., and Croteau, R. (1991) *Plant Physiol.* **96**:44-49; Funk, C., Lewinsohn, E., Stofer Vogel, B., Steele C., and Croteau, R. (1994) *Plant Physiol.* **106**:999-1005). Therefore, Grand fir has been developed as a model system to study the biochemical and molecular genetic regulation of constitutive and inducible terpene biosynthesis in conifers (Steele, C., Lewinsohn, E., and Croteau, R. (1995) *Proc. Natl. Acad. Sci. USA* **92**:4164-4168).

Most monoterpenes are derived from geranyl diphosphate, the ubiquitous C₁₀ intermediate of the isoprenoid pathway, by synthases which catalyze the divalent metal ion-dependent ionization (to 1, FIGURE 1) and isomerization of this substrate to enzyme-bound linalyl diphosphate which, following rotation about C2-C3, undergoes a second ionization (to 2, FIGURE 1) followed by cyclization to the α -terpinyl cation, the first cyclic intermediate en route to both monocyclic and bicyclic products (Croteau, R., and Cane, D.E. (1985) *Methods Enzymol.* **110**:383-405; Croteau, R. (1987) *Chem. Rev.* **87**:929-954) (FIGURE 1). Acyclic monoterpenes, such as myrcene, may arise by deprotonation of carbocations 1 or 2, whereas the isomerization step to linalyl diphosphate is required in the case of cyclic

types, such as limonene and pinenes, which cannot be derived from geranyl diphosphate directly because of the geometric impediment of the *trans*-double bond at C2-C3 (Croteau, R., and Cane, D.E. (1985) *Methods Enzymol.* **110**:383-405; Croteau, R. (1987) *Chem. Rev.* **87**:929-954). Many monoterpene synthases catalyze the formation of multiple products, including acyclic, monocyclic and bicyclic types, by variations on this basic mechanism (Gambliel, H., and Croteau, R. (1984) *J. Biol. Chem.* **259**:740-748; Croteau, R., Satterwhite, D.M., Cane, D.E., and Chang, C.C. (1988) *J. Biol. Chem.* **263**:10063-10071; Croteau, R., and Satterwhite, D.M. (1989) *J. Biol. Chem.* **264**:15309-15315). For example, (-)-limonene synthase, the principal monoterpene synthase of spearmint (*Mentha spicata*) and peppermint (*M. x piperita*), produces small amounts of myrcene, (-)- α -pinene and (-)- β -pinene in addition to the monocyclic product (Rajaonarivony, J.I.M., Gershenzon, J., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **296**:49-57; Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* **268**:23016-23024). Conversely, six different inducible monoterpene synthase activities have been demonstrated in extracts of wounded Grand fir stem (Gijzen, M., Lewinsohn, E., and Croteau, R. (1991) *Arch. Biochem. Biophys.* **289**:267-273) indicating that formation of acyclic, monocyclic and bicyclic monoterpenes in this species involves several genes encoding distinct catalysts. The inducible (-)-pinene synthase has been purified (Lewinsohn, E., Gijzen, M., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **293**:167-173), and isotopically sensitive branching experiments employed to demonstrate that this enzyme synthesizes both (-)- α - and (-)- β -pinene (Wagschal, K., Savage, T.J., and Croteau, R. (1991) *Tetrahedron* **47**:5933-5944).

Deciphering the molecular genetic control of oleoresinosis and examining structure-function relationships among the monoterpene synthases of Grand fir requires isolation of the cDNA species encoding these key enzymes. Although a protein-based cloning strategy was recently employed to acquire a cDNA for the major wound-inducible diterpene synthase from Grand fir, abietadiene synthase (Funk, C., Lewinsohn, E., Stofer Vogel, B., Steele C., and Croteau, R. (1994) *Plant Physiol.* **106**:999-1005; LaFever, R.E., Stofer Vogel, B., and Croteau, R. (1994) *Arch. Biochem. Biophys.* **313**:139-149; Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268), all attempts at the reverse genetic approach to cloning of Grand fir monoterpene synthases have failed (Steele, C., Lewinsohn, E., and Croteau, R. (1995) *Proc. Natl. Acad. Sci. USA* **92**:4164-4168). As an alternative, a similarity-based PCR strategy was developed (Steele, C.,

- Lewinsohn, E., and Croteau, R. (1995) *Proc. Natl. Acad. Sci. USA* 92:4164-4168) that employed sequence information from terpene synthases of angiosperm origin, namely a monoterpene synthase, (-)-4S-limonene synthase, from spearmint (*Mentha spicata*, Lamiaceae) (Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* 268:23016-23024), a sesquiterpene synthase, 5-*epi*-aristolochene synthase, from tobacco (*Nicotiana tabacum*, Solanaceae) (Facchini, P.J., and Chappell, J. (1992) *Proc. Natl. Acad. Sci. USA* 89:11088-11092), and a diterpene synthase, casbene synthase, from castor bean (*Ricinus communis*, Euphorbiaceae) (Mau, C.J.D., and West, C.A. (1994) *Proc. Natl. Acad. Sci. USA* 91:8497-8501).

Monoterpenes have significant potential for cancer prevention and treatment. Monoterpenes such as limonene, perillyl alcohol, carvone, geraniol and farnesol not only reduce tumor incidence and slow tumor proliferation, but have also been reported to cause regression of established solid tumors by initiating apoptosis (Mills J.J., Chari R.S., Boyer I.J., Gould M.N., Jirtle R.L., *Cancer Res.*, 55:979-983, 1995). Terpenes have activity against cancers such as mammary, colon, and prostate. Clinical trials are being pursued (Seachrist L, *J. NIH Res.* 8:43) in patients with various types of advanced cancers to validate the health benefits of dietary terpenes for humans. However, terpenes are present in Western diets at levels that are probably inadequate for any significant preventive health benefits. Daily supplementation of the diet with a terpene concentrate (10-20 g/day) would appear to be the most rational strategy for dietary therapy of diagnosed cases of cancer. This invention envisages the production of such nutritionally beneficial terpenes in vegetable oils consumed daily via the engineering of relevant genes from Grand fir into oil seed crop plants such as oil seed brassica (canola), soybean and corn.

Brief Description of the Drawings

The foregoing aspects and many of the attendant advantages of this invention will become more readily appreciated as the same becomes better understood by reference to the following detailed description, when taken in conjunction with the accompanying drawings, wherein:

FIGURE 1 is a schematic representation depicting the mechanism for the conversion of geranyl diphosphate to myrcene, (-)-limonene, β -phellandrene, (-)- α -pinene and (-)- β -pinene by monoterpene synthases from Grand fir. Formation of the monocyclic and bicyclic products requires preliminary isomerization of geranyl diphosphate to linalyl diphosphate. The acyclic product could be formed

from either geranyl diphosphate or linalyl diphosphate via carbocations 1 or 2. *OPP* denotes the diphosphate moiety.

FIGURE 2 is a sequence comparison of plant terpene synthases. A three-letter designation (*Tps*) for the gene family is proposed with sub-groups (*Tpsa* through *Tpsf*) defined by a minimum of 40 % amino acid identity between members. The numbers in parenthesis are the references to the published sequences.

FIGURE 3 depicts a GLC-MS analysis of the products of the recombinant protein encoded by *AG2.2* (SEQ ID NO:1), the sequence of the protein encoded by clone *AG2.2* being set forth in SEQ ID NO:2. The GLC profile of the total pentane-soluble products generated from geranyl diphosphate when incubated with a cell-free extract of *E. coli* XL1-Blue/pGAG2.2 is illustrated (a), as are the mass fragmentation patterns for the monoterpene product with $R_t = 12.22$ min (b) and for authentic myrcene (c).

FIGURE 4 depicts a GLC-MS analysis of the products of the recombinant protein encoded by *AG3.18* (SEQ ID NO:3), the sequence of the protein encoded by clone *AG3.18* (SEQ ID NO:3) being set forth in SEQ ID NO:4. The GLC profile of the total pentane-soluble products generated from geranyl diphosphate when incubated with a cell-free extract of *E. coli* XL1-Blue/pGAG3.18 is illustrated (a), as are the mass fragmentation patterns (selected ion mode) for the monoterpene products with $R_t = 11.34$ min (b), and $R_t = 13.37$ min (d), and for authentic α -pinene (c) and authentic β -pinene (e).

FIGURE 5 depicts a GLC-MS analysis of the products of the recombinant protein encoded by *AG10* (SEQ ID NO:5), the sequence of the protein encoded by clone *AG10* (SEQ ID NO:5) being set forth in SEQ ID NO:6. The GLC profile of the total pentane-soluble products generated from geranyl diphosphate when incubated with a cell-free extract of *E. coli* BL21(DE3)/pSBAG10 is illustrated (a), as are the mass fragmentation patterns for the principal monoterpene product with $R_t = 13.93$ min (b) and for authentic limonene (c).

Summary of the Invention

In accordance with the foregoing, cDNAs encoding myrcene synthase, (-)-limonene synthase and (-)-pinene synthase from Grand fir (*Abies grandis*) have been isolated and sequenced, and the corresponding amino acid sequences have been deduced. Accordingly, the present invention relates to isolated DNA sequences which code for the expression of myrcene synthase, such as the sequence designated SEQ ID NO:1 which encodes myrcene synthase from Grand fir (*Abies grandis*), for

the expression of (-)-pinene synthase, such as the sequence designated SEQ ID NO:3, which encodes the (-)-pinene synthase from Grand fir (*Abies grandis*) and for the expression of (-)-limonene synthase, such as the sequence designated SEQ ID NO:5, which encodes (-)-limonene synthase from Grand fir (*Abies grandis*). In other aspects, the present invention is directed to replicable recombinant cloning vehicles comprising a nucleic acid sequence, e.g., a DNA sequence which codes for a myrcene synthase, (-)-limonene synthase or (-)-pinene synthase, or for a base sequence sufficiently complementary to at least a portion of DNA or RNA encoding myrcene synthase, (-)-limonene synthase or (-)-pinene synthase to enable hybridization therewith (e.g., antisense RNA or fragments of DNA complementary to a portion of DNA or RNA molecules encoding myrcene synthase, (-)-limonene synthase or (-)-pinene synthase which are useful as polymerase chain reaction primers or as probes for any of the foregoing synthases or related genes). In yet other aspects of the invention, modified host cells are provided that have been transformed, transfected, infected and/or injected with a recombinant cloning vehicle and/or DNA sequence of the invention. Thus, the present invention provides for the recombinant expression of myrcene synthase, (-)-limonene synthase and (-)-pinene synthase, and the inventive concepts may be used to facilitate the production, isolation and purification of significant quantities of recombinant myrcene synthase, (-)-limonene synthase and (-)-pinene synthase (or of their primary enzyme products) for subsequent use, to obtain expression or enhanced expression of myrcene synthase, (-)-limonene synthase and (-)-pinene synthase in plants, microorganisms or animals, or may be otherwise employed in an environment where the regulation or expression of myrcene synthase, (-)-limonene synthase and (-)-pinene synthase is desired for the production of these synthases, or their enzyme products, or derivatives thereof. In another aspect, the present invention relates to manipulation of monoterpene production to enhance resistance to insects and/or accumulate nutritionally beneficial monoterpenes in oil seeds.

Detailed Description of the Preferred Embodiment

As used herein, the terms "amino acid" and "amino acids" refer to all naturally occurring L- α -amino acids or their residues. The amino acids are identified by either the single-letter or three-letter designations:

Asp	D	aspartic acid	Ile	I	isoleucine
Thr	T	threonine	Leu	L	leucine
Ser	S	serine	Tyr	Y	tyrosine

	Glu	E	glutamic acid	Phe	F	phenylalanine
	Pro	P	proline	His	H	histidine
	Gly	G	glycine	Lys	K	lysine
	Ala	A	alanine	Arg	R	arginine
5	Cys	C	cysteine	Trp	W	tryptophan
	Val	V	valine	Gln	Q	glutamine
	Met	M	methionine	Asn	N	asparagine

As used herein, the term "nucleotide" means a monomeric unit of DNA or RNA containing a sugar moiety (pentose), a phosphate and a nitrogenous heterocyclic base. The base is linked to the sugar moiety via the glycosidic carbon (1' carbon of pentose) and that combination of base and sugar is called a nucleoside. The base characterizes the nucleotide with the four bases of DNA being adenine ("A"), guanine ("G"), cytosine ("C") and thymine ("T"). Inosine ("I") is a synthetic base that can be used to substitute for any of the four, naturally-occurring bases (A, C, G or T). The four RNA bases are A, G, C and uracil ("U"). The nucleotide sequences described herein comprise a linear array of nucleotides connected by phosphodiester bonds between the 3' and 5' carbons of adjacent pentoses.

The term "percent identity" means the percentage of amino acids or nucleotides that occupy the same relative position when two amino acid sequences, or two nucleic acid sequences are aligned side by side.

The term "percent similarity" is a statistical measure of the degree of relatedness of two compared protein sequences. The percent similarity is calculated by a computer program that assigns a numerical value to each compared pair of amino acids based on chemical similarity (e.g., whether the compared amino acids are acidic, basic, hydrophobic, aromatic, etc.) and/or evolutionary distance as measured by the minimum number of base pair changes that would be required to convert a codon encoding one member of a pair of compared amino acids to a codon encoding the other member of the pair. Calculations are made after a best fit alignment of the two sequences have been made empirically by iterative comparison of all possible alignments. (Henikoff, S. and Henikoff, J.G., *Proc. Nat'l. Acad. Sci. USA* **89**:10915-10919, 1992).

"Oligonucleotide" refers to short length single or double stranded sequences of deoxyribonucleotides linked via phosphodiester bonds. The oligonucleotides are chemically synthesized by known methods and purified, for example, on polyacrylamide gels.

The term "myrcene synthase" is used herein to mean an enzyme capable of generating multiple monoterpenes from geranyl diphosphate. The principal and characteristic monoterpene synthesized by myrcene synthase is myrcene, which constitutes at least about 60% of the monoterpene mixture synthesized by myrcene synthase from geranyl diphosphate.

The term "(-)-limonene synthase" is used herein to mean an enzyme capable of generating multiple monoterpenes from geranyl diphosphate. The principal and characteristic monoterpene synthesized by (-)-limonene synthase is (-)-limonene, which constitutes at least about 60% of the monoterpene mixture synthesized by (-)-limonene synthase from geranyl diphosphate.

The term "(-)-pinene synthase" is used herein to mean an enzyme capable of generating multiple monoterpenes from geranyl diphosphate. The principal and characteristic monoterpene synthesized by (-)-pinene synthase is (-)-pinene, which comprises at least about 60% of the monoterpene mixture synthesized by (-)-pinene synthase from geranyl diphosphate.

The abbreviation "SSPE" refers to a buffer used in nucleic acid hybridization solutions. The 20X (twenty times concentrate) stock SSPE buffer solution is prepared as follows: dissolve 175.3 grams of NaCl, 27.6 grams of $\text{NaH}_2\text{PO}_4\text{H}_2\text{O}$ and 7.4 grams of EDTA in 800 millilitres of H_2O . Adjust the pH to pH 7.4 with NaOH. Adjust the volume to one liter with H_2O .

The terms "alteration", "amino acid sequence alteration", "variant" and "amino acid sequence variant" refer to monoterpene synthase molecules with some differences in their amino acid sequences as compared to the corresponding, native, i.e., naturally-occurring, monoterpene synthases. Ordinarily, the variants will possess at least about 70% homology with the corresponding native monoterpene synthases, and preferably, they will be at least about 80% homologous with the corresponding, native monoterpene synthases. The amino acid sequence variants of the monoterpene synthases falling within this invention possess substitutions, deletions, and/or insertions at certain positions. Sequence variants of monoterpene synthases may be used to attain desired enhanced or reduced enzymatic activity, modified regiochemistry or stereochemistry, or altered substrate utilization or product distribution.

Substitutional monoterpene synthase variants are those that have at least one amino acid residue in the native monoterpene synthase sequence removed and a different amino acid inserted in its place at the same position. The substitutions may

be single, where only one amino acid in the molecule has been substituted, or they may be multiple, where two or more amino acids have been substituted in the same molecule. Substantial changes in the activity of the monoterpene synthase molecules of the present invention may be obtained by substituting an amino acid with a side chain that is significantly different in charge and/or structure from that of the native amino acid. This type of substitution would be expected to affect the structure of the polypeptide backbone and/or the charge or hydrophobicity of the molecule in the area of the substitution.

Moderate changes in the activity of the monoterpene synthase molecules of the present invention would be expected by substituting an amino acid with a side chain that is similar in charge and/or structure to that of the native molecule. This type of substitution, referred to as a conservative substitution, would not be expected to substantially alter either the structure of the polypeptide backbone or the charge or hydrophobicity of the molecule in the area of the substitution.

Insertional monoterpene synthase variants are those with one or more amino acids inserted immediately adjacent to an amino acid at a particular position in the native monoterpene synthase molecule. Immediately adjacent to an amino acid means connected to either the α -carboxy or α -amino functional group of the amino acid. The insertion may be one or more amino acids. Ordinarily, the insertion will consist of one or two conservative amino acids. Amino acids similar in charge and/or structure to the amino acids adjacent to the site of insertion are defined as conservative. Alternatively, this invention includes insertion of an amino acid with a charge and/or structure that is substantially different from the amino acids adjacent to the site of insertion.

Deletional variants are those where one or more amino acids in the native monoterpene synthase molecules have been removed. Ordinarily, deletional variants will have one or two amino acids deleted in a particular region of the monoterpene synthase molecule.

The terms "biological activity", "biologically active", "activity" and "active" refer to the ability of the monoterpene synthases of the present invention to convert geranyl diphosphate to a group of monoterpenes, of which myrcene is the principal and characteristic monoterpene synthesized by myrcene synthase, (-)-limonene is the principal and characteristic monoterpene synthesized by (-)-limonene synthase and (-)-pinene is the principal and characteristic monoterpene synthesized by (-)-pinene synthase. The monoterpenes produced by the monoterpene synthases of the present

invention are as measured in an enzyme activity assay, such as the assay described in Example 3. Amino acid sequence variants of the terpene synthases of the present invention may have desirable altered biological activity including, for example, altered reaction kinetics, substrate utilization product distribution or other characteristics such as regiochemistry and stereochemistry.

The terms "DNA sequence encoding", "DNA encoding" and "nucleic acid encoding" refer to the order or sequence of deoxyribonucleotides along a strand of deoxyribonucleic acid. The order of these deoxyribonucleotides determines the order of amino acids along the translated polypeptide chain. The DNA sequence thus codes for the amino acid sequence.

The terms "replicable expression vector" and "expression vector" refer to a piece of DNA, usually double-stranded, which may have inserted into it a piece of foreign DNA. Foreign DNA is defined as heterologous DNA, which is DNA not naturally found in the host. The vector is used to transport the foreign or heterologous DNA into a suitable host cell. Once in the host cell, the vector can replicate independently of or coincidental with the host chromosomal DNA, and several copies of the vector and its inserted (foreign) DNA may be generated. In addition, the vector contains the necessary elements that permit translating the foreign DNA into a polypeptide. Many molecules of the polypeptide encoded by the foreign DNA can thus be rapidly synthesized.

The terms "transformed host cell," "transformed" and "transformation" refer to the introduction of DNA into a cell. The cell is termed a "host cell", and it may be a prokaryotic or a eukaryotic cell. Typical prokaryotic host cells include various strains of *E. coli*. Typical eukaryotic host cells are plant cells, such as maize cells, yeast cells, insect cells or animal cells. The introduced DNA is usually in the form of a vector containing an inserted piece of DNA. The introduced DNA sequence may be from the same species as the host cell or from a different species from the host cell, or it may be a hybrid DNA sequence, containing some foreign DNA and some DNA derived from the host species.

The following abbreviations are used herein: bp(s), base pair(s); DEAE, *O*-diethylaminoethyl; DTT, dithiothreitol; EDTA, ethylenediaminetetraacetic acid; GC, gas chromatography; IPTG, isopropyl- β -D-thiogalactopyranoside; LB, Luria-Bertani; Mopso, 3-(*N*-morpholino)-2-hydroxypropane-sulfonic acid; MS, mass spectrum/ spectrometry; nt(s), nucleotide(s); ORF, open reading frame; PCR,

polymerase chain reaction; PVDF, polyvinylidenedifluoride; SDS, sodium dodecyl sulfate; Tris, tris(hydroxymethyl) aminomethane; UV, ultraviolet.

In accordance with the present invention, cDNAs encoding myrcene synthase (SEQ ID NO:1), (-)-pinene synthase (SEQ ID NO:3) and (-)-limonene synthase (SEQ ID NO:5) from Grand fir (*Abies grandis*) were isolated and sequenced in the following manner. Based on comparison of sequences of limonene synthase from spearmint (Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* **268**:23016-23024), 5-*epi*-aristolochene synthase from tobacco (Facchini, P.J., and Chappell, J. (1992) *Proc. Natl. Acad. Sci. USA* **89**:11088-11092), and casbene synthase from castor bean (Mau, C.J.D., and West, C.A. (1994) *Proc. Natl. Acad. Sci. USA* **91**:8497-8501), four conserved regions were identified for which a set of consensus, degenerate primers were synthesized: Primer A (SEQ ID NO:7), Primer B (SEQ ID NO:8), Primer C (SEQ ID NO:9) and Primer D (SEQ ID NO:10). Primers A (SEQ ID NO:7), B (SEQ ID NO:8), and D (SEQ ID NO:10) were sense primers, while Primer C (SEQ ID NO:9), was an antisense primer. Each of the sense primers, A (SEQ ID NO:7), B (SEQ ID NO:8) and D (SEQ ID NO:10), was used for PCR in combination with antisense primer C (SEQ ID NO:9) by employing a broad range of amplification conditions. Analysis of the PCR reaction products by agarose gel electrophoresis revealed that only the combination of primers C (SEQ ID NO:9) and D (SEQ ID NO:10) generated a specific PCR product of approximately 110 bps.

The 110 bps PCR product was gel purified, ligated into a plasmid, and transformed into *E. coli* XL1-Blue cells. Plasmid DNA was prepared from 41 individual transformants and the inserts were sequenced. Four different insert sequences were identified, and were designated as probes 1 (SEQ ID NO:11), 2 (SEQ ID NO:12), 4 (SEQ ID NO:13) and 5 (SEQ ID NO:14). Probes 1 (SEQ ID NO:11), 2 (SEQ ID NO:12), 4 (SEQ ID NO:13) and 5 (SEQ ID NO:14) were used to screen a cDNA library made from mRNA extracted from wounded Grand fir stems, and the longest clone that hybridized to each of these probes was isolated and sequenced. Thus, clone *AG1.28* (SEQ ID NO:15) is the longest cDNA clone that hybridized to probe 1 (SEQ ID NO:11), clone *AG2.2* (SEQ ID NO:1) is the longest cDNA clone that hybridized to probe 2 (SEQ ID NO:12), clone *AG4.30* (SEQ ID NO:17) is the longest cDNA clone that hybridized to probe 4 (SEQ ID NO: 13), and clone *AG5.9* (SEQ ID NO:19) is the longest cDNA clone that hybridized to probe 5 (SEQ ID NO:14).

Truncated clone *AG1.28* (SEQ ID NO:15) resembled most closely in size and sequence (72% similarity, 49% identity) a diterpene cyclase, abietadiene synthase, from Grand fir. Clones *AG4.30* (SEQ ID NO:17) and *AG5.9* (SEQ ID NO:19) encode sesquiterpene synthases. Sequence and functional analysis of clone *AG2.2* (SEQ ID NO:1) revealed that it encoded the monoterpene synthase, myrcene synthase.

Alignment of the four new terpene synthase cDNA sequences *AG1.28* (SEQ ID NO:15), *AG2.2* (SEQ ID NO:1), *AG4.30* (SEQ ID NO:17) and *AG5.9* (SEQ ID NO:19) with that for abietadiene synthase (Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268) allowed the identification of several, conserved sequence motifs. Two new sense PCR primers, primer E (SEQ ID NO:21) and primer F (SEQ ID NO:22) were designed based on the sequence of the conserved protein sequence motifs. A new antisense PCR primer, primer G (SEQ ID NO:23), was designed based on limited sequence information available from pinene synthase. The combination of primer E (SEQ ID NO:21) and primer G (SEQ ID NO:23) amplified a cDNA product of 1022 bps, which was designated as probe 3 (SEQ ID NO:24).

Probe 3 (SEQ ID NO:24) was used to screen a cDNA library made from mRNA extracted from wounded Grand fir stems. Hybridization of 10^5 Grand fir λ ZAP II cDNA clones with probe 3 (SEQ ID NO:24) yielded two types of signals comprised of about 400 strongly positive clones and an equal number of weak positives, indicating that the probe recognized more than one type of cDNA. Thirty-four of the former clones and eighteen of the latter were purified, the inserts were selected by size (2.0-2.5 kb), and the *in vivo* excised clones were partially sequenced from both ends. Those clones which afforded weak hybridization signals were shown to contain inserts that were either identical to myrcene synthase clone *AG2.2* (SEQ ID NO:1) or exhibited no significant sequence similarity to terpene synthases.

Clones which gave strong hybridization signals segregated into distinct sequence groups represented by clone *AG3.18* (SEQ ID NO:3) and clone *AG10* (SEQ ID NO:5). Both *AG3.18* (SEQ ID NO:3) and *AG10* (SEQ ID NO:5) were subcloned into plasmid expression vectors and expressed in *E. coli*. When extracts of the induced cells were tested for terpene synthase activity with all of the potential prenyl diphosphate substrates, only geranyl diphosphate was utilized. Extracts from *E. coli* containing the *AG10* (SEQ ID NO:5) expression construct converted geranyl diphosphate to the -4S enantiomer of limonene as the major product, indicating that

5 *AG10* (SEQ ID NO:5) encodes (-)-limonene synthase. Similar analysis of the monoterpene products generated from geranyl diphosphate by cell-free extracts of *E. coli* containing the *AG3.18* (SEQ ID NO:3) insert ligated into an expression vector revealed the presence of a 42:58% mixture of α -pinene and β -pinene, the same product ratio previously described for the purified, native (-)-pinene synthase from Grand fir. These data indicate that *AG3.18* (SEQ ID NO:3) encodes (-)-pinene synthase.

10 The isolation of cDNAs encoding (-)-limonene synthase, (-)-pinene synthase and myrcene synthase permits the development of efficient expression systems for these functional enzymes; provides useful tools for examining the developmental regulation of monoterpene biosynthesis; permits investigation of the reaction mechanism(s) of these unusual, multiproduct enzymes, and permits the isolation of other (-)-limonene synthase, (-)-pinene synthase and myrcene synthases. The isolation of the (-)-limonene synthase, (-)-pinene synthase and myrcene synthase cDNAs also permits the transformation of a wide range of organisms in order to introduce monoterpene biosynthesis *de novo*, or to modify endogenous monoterpene biosynthesis.

20 Although the (-)-limonene synthase, (-)-pinene synthase and myrcene synthase proteins set forth in SEQ ID NO:6, SEQ ID NO:4 and SEQ ID NO:2, respectively, direct the enzymes to plastids, substitution of the presumptive targeting sequence of each of these enzymes (SEQ ID NO:2, amino acids 1 to 61; SEQ ID NO:4, amino acids 1 to 61; SEQ ID NO:6, amino acids 1 to 66) with other transport sequences well known in the art (see, e.g., von Heijne et al., *Eur. J. Biochem.* **180**:535-545, 1989; Stryer, *Biochemistry*, W.H. Freeman and Company, New York, NY, p. 769 [1988]) may be employed to direct the (-)-limonene synthase, (-)-pinene synthase and myrcene synthase to other cellular or extracellular locations.

30 In addition to the native (-)-limonene synthase, (-)-pinene synthase and myrcene synthase amino acid sequences of SEQ ID NO:6, SEQ ID NO:4 and SEQ ID NO:2, respectively, sequence variants produced by deletions, substitutions, mutations and/or insertions are intended to be within the scope of the invention except insofar as limited by the prior art. The (-)-limonene synthase, (-)-pinene synthase and myrcene synthase amino acid sequence variants of this invention may be constructed by mutating the DNA sequences that encode the wild-type synthases, such as by using techniques commonly referred to as site-directed mutagenesis.

35 Nucleic acid molecules encoding the monoterpene synthases of the present invention

can be mutated by a variety of PCR techniques well known to one of ordinary skill in the art. See, e.g., "PCR Strategies", M.A. Innis, D.H. Gelfand and J.J. Sninsky, eds., 1995, Academic Press, San Diego, CA (Chapter 14); "PCR Protocols: A Guide to Methods and Applications", M.A. Innis, D.H. Gelfand, J.J. Sninsky and T.J. White, eds., Academic Press, NY (1990).

By way of non-limiting example, the two primer system utilized in the Transformer Site-Directed Mutagenesis kit from Clontech, may be employed for introducing site-directed mutants into the monoterpene synthase genes of the present invention. Following denaturation of the target plasmid in this system, two primers are simultaneously annealed to the plasmid; one of these primers contains the desired site-directed mutation, the other contains a mutation at another point in the plasmid resulting in elimination of a restriction site. Second strand synthesis is then carried out, tightly linking these two mutations, and the resulting plasmids are transformed into a *mutS* strain of *E. coli*. Plasmid DNA is isolated from the transformed bacteria, restricted with the relevant restriction enzyme (thereby linearizing the unmutated plasmids), and then retransformed into *E. coli*. This system allows for generation of mutations directly in an expression plasmid, without the necessity of subcloning or generation of single-stranded phagemids. The tight linkage of the two mutations and the subsequent linearization of unmutated plasmids results in high mutation efficiency and allows minimal screening. Following synthesis of the initial restriction site primer, this method requires the use of only one new primer type per mutation site. Rather than prepare each positional mutant separately, a set of "designed degenerate" oligonucleotide primers can be synthesized in order to introduce all of the desired mutations at a given site simultaneously. Transformants can be screened by sequencing the plasmid DNA through the mutagenized region to identify and sort mutant clones. Each mutant DNA can then be restricted and analyzed by electrophoresis on Mutation Detection Enhancement gel (J.T. Baker) to confirm that no other alterations in the sequence have occurred (by band shift comparison to the unmutagenized control).

The verified mutant duplexes in the pET (or other) overexpression vector can be employed to transform *E. coli* such as strain *E. coli* BL21(DE3)pLysS, for high level production of the mutant protein, and purification by standard protocols. The method of FAB-MS mapping can be employed to rapidly check the fidelity of mutant expression. This technique provides for sequencing segments throughout the whole protein and provides the necessary confidence in the sequence assignment. In a

mapping experiment of this type, protein is digested with a protease (the choice will depend on the specific region to be modified since this segment is of prime interest and the remaining map should be identical to the map of unmutagenized protein). The set of cleavage fragments is fractionated by microbore HPLC (reversed phase or ion exchange, again depending on the specific region to be modified) to provide several peptides in each fraction, and the molecular weights of the peptides are determined by FAB-MS. The masses are then compared to the molecular weights of peptides expected from the digestion of the predicted sequence, and the correctness of the sequence quickly ascertained. Since this mutagenesis approach to protein modification is directed, sequencing of the altered peptide should not be necessary if the MS agrees with prediction. If necessary to verify a changed residue, CAD-tandem MS/MS can be employed to sequence the peptides of the mixture in question, or the target peptide purified for subtractive Edman degradation or carboxypeptidase Y digestion depending on the location of the modification.

In the design of a particular site directed mutagenesis, it is generally desirable to first make a non-conservative substitution (e.g., Ala for Cys, His or Glu) and determine if activity is greatly impaired as a consequence. The properties of the mutagenized protein are then examined with particular attention to the kinetic parameters of K_m and k_{cat} as sensitive indicators of altered function, from which changes in binding and/or catalysis *per se* may be deduced by comparison to the native enzyme. If the residue is by this means demonstrated to be important by activity impairment, or knockout, then conservative substitutions can be made, such as Asp for Glu to alter side chain length, Ser for Cys, or Arg for His. For hydrophobic segments, it is largely size that is usefully altered, although aromatics can also be substituted for alkyl side chains. Changes in the normal product distribution can indicate which step(s) of the reaction sequence have been altered by the mutation. Modification of the hydrophobic pocket can be employed to change binding conformations for substrates and result in altered regiochemistry and/or stereochemistry.

Other site directed mutagenesis techniques may also be employed with the nucleotide sequences of the invention. For example, restriction endonuclease digestion of DNA followed by ligation may be used to generate deletion variants of (-)-limonene synthase, (-)-pinene synthase and myrcene synthase, as described in section 15.3 of Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, New York, NY [1989]). A similar strategy

may be used to construct insertion variants, as described in section 15.3 of Sambrook et al., *supra*.

Oligonucleotide-directed mutagenesis may also be employed for preparing substitution variants of this invention. It may also be used to conveniently prepare the deletion and insertion variants of this invention. This technique is well known in the art as described by Adelman et al. (*DNA* 2:183 [1983]); Sambrook et al., *supra*; "Current Protocols in Molecular Biology", 1991, Wiley (NY), F.T. Ausubel, R. Brent, R.E. Kingston, D.D. Moore, J.D. Seidman, J.A. Smith and K. Struhl, eds.

Generally, oligonucleotides of at least 25 nucleotides in length are used to insert, delete or substitute two or more nucleotides in the (-)-limonene synthase, (-)-pinene synthase and myrcene synthase molecule. An optimal oligonucleotide will have 12 to 15 perfectly matched nucleotides on either side of the nucleotides coding for the mutation. To mutagenize wild-type (-)-limonene synthase, (-)-pinene synthase and myrcene synthase, the oligonucleotide is annealed to the single-stranded DNA template molecule under suitable hybridization conditions. A DNA polymerizing enzyme, usually the Klenow fragment of *E. coli* DNA polymerase I, is then added. This enzyme uses the oligonucleotide as a primer to complete the synthesis of the mutation-bearing strand of DNA. Thus, a heteroduplex molecule is formed such that one strand of DNA encodes the wild-type synthase inserted in the vector, and the second strand of DNA encodes the mutated form of the synthase inserted into the same vector. This heteroduplex molecule is then transformed into a suitable host cell.

Mutants with more than one amino acid substituted may be generated in one of several ways. If the amino acids are located close together in the polypeptide chain, they may be mutated simultaneously using one oligonucleotide that codes for all of the desired amino acid substitutions. If however, the amino acids are located some distance from each other (separated by more than ten amino acids, for example) it is more difficult to generate a single oligonucleotide that encodes all of the desired changes. Instead, one of two alternative methods may be employed. In the first method, a separate oligonucleotide is generated for each amino acid to be substituted. The oligonucleotides are then annealed to the single-stranded template DNA simultaneously, and the second strand of DNA that is synthesized from the template will encode all of the desired amino acid substitutions. An alternative method involves two or more rounds of mutagenesis to produce the desired mutant. The first round is as described for the single mutants: wild-type (-)-limonene synthase,

(-)-pinene synthase and myrcene synthase DNA is used for the template, an oligonucleotide encoding the first desired amino acid substitution(s) is annealed to this template, and the heteroduplex DNA molecule is then generated. The second round of mutagenesis utilizes the mutated DNA produced in the first round of mutagenesis as the template. Thus, this template already contains one or more mutations. The oligonucleotide encoding the additional desired amino acid substitution(s) is then annealed to this template, and the resulting strand of DNA now encodes mutations from both the first and second rounds of mutagenesis. This resultant DNA can be used as a template in a third round of mutagenesis, and so on.

A gene encoding (-)-limonene synthase, (-)-pinene synthase and myrcene synthase may be incorporated into any organism (intact plant, animal, microbe, etc.), or cell culture derived therefrom, that produces geranyl diphosphate. A (-)-limonene synthase, (-)-pinene synthase and myrcene synthase gene may be introduced into any organism for a variety of purposes including, but not limited to: production of (-)-limonene synthase, (-)-pinene synthase and myrcene synthase, or their products; production or modification of flavor and aroma properties; improvement of defense capability, and the alteration of other ecological interactions mediated by myrcene, (-)-limonene, (-)-pinene, or their derivatives.

Eukaryotic expression systems may be utilized for the production of (-)-limonene synthase, (-)-pinene synthase and myrcene synthase since they are capable of carrying out any required posttranslational modifications and of directing the enzymes to the proper membrane location. A representative eukaryotic expression system for this purpose uses the recombinant baculovirus, *Autographa californica* nuclear polyhedrosis virus (AcNPV; M.D. Summers and G.E. Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures* [1986]; Luckow et al., *Bio-technology* 6:47-55 [1987]) for expression of the terpenoid synthases of the invention. Infection of insect cells (such as cells of the species *Spodoptera frugiperda*) with the recombinant baculoviruses allows for the production of large amounts of the monoterpenoid synthase proteins. In addition, the baculovirus system has other important advantages for the production of recombinant (-)-limonene synthase, (-)-pinene synthase and myrcene synthase. For example, baculoviruses do not infect humans and can therefore be safely handled in large quantities. In the baculovirus system, a DNA construct is prepared including a DNA segment encoding (-)-limonene synthase, (-)-pinene synthase and myrcene synthase and a vector. The vector may comprise the polyhedron gene promoter region of a

5 baculovirus, the baculovirus flanking sequences necessary for proper cross-over during recombination (the flanking sequences comprise about 200-300 base pairs adjacent to the promoter sequence) and a bacterial origin of replication which permits the construct to replicate in bacteria. The vector is constructed so that (i) the DNA segment is placed adjacent (or operably linked or "downstream" or "under the control of") to the polyhedron gene promoter and (ii) the promoter/monoterpene synthase combination is flanked on both sides by 200-300 base pairs of baculovirus DNA (the flanking sequences).

10 To produce the monoterpene synthase DNA construct, a cDNA clone encoding the full length (-)-limonene synthase, (-)-pinene synthase and myrcene synthase is obtained using methods such as those described herein. The DNA construct is contacted in a host cell with baculovirus DNA of an appropriate baculovirus (that is, of the same species of baculovirus as the promoter encoded in the construct) under conditions such that recombination is effected. The resulting
15 recombinant baculoviruses encode the full (-)-limonene synthase, (-)-pinene synthase and myrcene synthase. For example, an insect host cell can be cotransfected or transfected separately with the DNA construct and a functional baculovirus. Resulting recombinant baculoviruses can then be isolated and used to infect cells to effect production of the monoterpene synthase. Host insect cells include, for
20 example, *Spodoptera frugiperda* cells, that are capable of producing a baculovirus-expressed monoterpene synthase. Insect host cells infected with a recombinant baculovirus of the present invention are then cultured under conditions allowing expression of the baculovirus-encoded (-)-limonene synthase, (-)-pinene synthase and myrcene synthase. (-)-limonene synthase, (-)-pinene synthase and myrcene synthase
25 thus produced are then extracted from the cells using methods known in the art.

Other eukaryotic microbes such as yeasts may also be used to practice this invention. The baker's yeast *Saccharomyces cerevisiae*, is a commonly used yeast, although several other strains are available. The plasmid YRp7 (Stinchcomb et al., *Nature* **282**:39 [1979]; Kingsman et al., *Gene* **7**:141 [1979]; Tschemper et al., *Gene*
30 **10**:157 [1980]) is commonly used as an expression vector in *Saccharomyces*. This plasmid contains the *trp1* gene that provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, such as strains ATCC No. 44,076 and PEP4-1 (Jones, *Genetics* **85**:12 [1977]). The presence of the *trp1* lesion as a characteristic of the yeast host cell genome then provides an effective environment
35 for detecting transformation by growth in the absence of tryptophan. Yeast host cells

are generally transformed using the polyethylene glycol method, as described by Hinnen (*Proc. Natl. Acad. Sci. USA* **75**:1929 [1978]). Additional yeast transformation protocols are set forth in Gietz et al., *N.A.R.* **20**(17) 1425(1992); Reeves et al., *FEMS* **99**(2-3):193-197, (1992).

5 Suitable promoting sequences in yeast vectors include the promoters for 3-phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* **255**:2073 [1980]) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* **7**:149 [1968]; Holland et al., *Biochemistry* **17**:4900 [1978]), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, 10 glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. In the construction of suitable expression plasmids, the termination sequences associated with these genes are also ligated into the expression vector 3' of the sequence desired to be expressed to provide polyadenylation of the mRNA and termination. Other 15 promoters that have the additional advantage of transcription controlled by growth conditions are the promoter region for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, and the aforementioned glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Any plasmid vector containing 20 yeast-compatible promoter, origin of replication and termination sequences is suitable.

Cell cultures derived from multicellular organisms, such as plants, may be used as hosts to practice this invention. Transgenic plants can be obtained, for example, by transferring plasmids that encode (-)-limonene synthase, (-)-pinene 25 synthase and myrcene synthase and a selectable marker gene, e.g., the kan gene encoding resistance to kanamycin, into *Agrobacterium tumifaciens* containing a helper Ti plasmid as described in Hoeckema et al., *Nature* **303**:179-181 [1983] and culturing the *Agrobacterium* cells with leaf slices of the plant to be transformed as described by An et al., *Plant Physiology* **81**:301-305 [1986]. Transformation of 30 cultured plant host cells is normally accomplished through *Agrobacterium tumifaciens*, as described above. Cultures of mammalian host cells and other host cells that do not have rigid cell membrane barriers are usually transformed using the calcium phosphate method as originally described by Graham and Van der Eb (*Virology* **52**:546 [1978]) and modified as described in sections 16.32-16.37 of 35 Sambrook et al., *supra*. However, other methods for introducing DNA into cells

such as Polybrene (Kawai and Nishizawa, *Mol. Cell. Biol.* 4:1172 [1984]), protoplast fusion (Schaffner, *Proc. Natl. Acad. Sci. USA* 77:2163 [1980]), electroporation (Neumann et al., *EMBO J.* 1:841 [1982]), and direct microinjection into nuclei (Capecchi, *Cell* 22:479 [1980]) may also be used. Additionally, animal transformation strategies are reviewed in Monastersky G.M. and Robl, J.M., *Strategies in Transgenic Animal Science*, ASM Press, Washington, D.C., 1995. Transformed plant calli may be selected through the selectable marker by growing the cells on a medium containing, e.g., kanamycin, and appropriate amounts of phytohormone such as naphthalene acetic acid and benzyladenine for callus and shoot induction. The plant cells may then be regenerated and the resulting plants transferred to soil using techniques well known to those skilled in the art.

In addition, a gene regulating (-)-limonene synthase, (-)-pinene synthase and myrcene synthase production can be incorporated into the plant along with a necessary promoter which is inducible. In the practice of this embodiment of the invention, a promoter that only responds to a specific external or internal stimulus is fused to the target cDNA. Thus, the gene will not be transcribed except in response to the specific stimulus. As long as the gene is not being transcribed, its gene product is not produced.

An illustrative example of a responsive promoter system that can be used in the practice of this invention is the glutathione-S-transferase (GST) system in maize. GSTs are a family of enzymes that can detoxify a number of hydrophobic electrophilic compounds that often are used as pre-emergent herbicides (Weigand et al., *Plant Molecular Biology* 7:235-243 [1986]). Studies have shown that the GSTs are directly involved in causing this enhanced herbicide tolerance. This action is primarily mediated through a specific 1.1 kb mRNA transcription product. In short, maize has a naturally occurring quiescent gene already present that can respond to external stimuli and that can be induced to produce a gene product. This gene has previously been identified and cloned. Thus, in one embodiment of this invention, the promoter is removed from the GST responsive gene and attached to a (-)-limonene synthase, (-)-pinene synthase and myrcene synthase gene that previously has had its native promoter removed. This engineered gene is the combination of a promoter that responds to an external chemical stimulus and a gene responsible for successful production of (-)-limonene synthase, (-)-pinene synthase and myrcene synthase.

In addition to the methods described above, several methods are known in the art for transferring cloned DNA into a wide variety of plant species, including gymnosperms, angiosperms, monocots and dicots (see, e.g., Glick and Thompson, eds., *Methods in Plant Molecular Biology*, CRC Press, Boca Raton, Florida [1993]). Representative examples include electroporation-facilitated DNA uptake by protoplasts (Rhodes et al., *Science* **240(4849)**:204-207 [1988]); treatment of protoplasts with polyethylene glycol (Lyznik et al., *Plant Molecular Biology* **13**:151-161 [1989]); and bombardment of cells with DNA laden microprojectiles (Klein et al., *Plant Physiol.* **91**:440-444 [1989] and Boynton et al., *Science* **240(4858)**:1534-1538 [1988]). Additionally, plant transformation strategies and techniques are reviewed in Birch, R.G., *Ann Rev Plant Phys Plant Mol Biol* **48**:297 (1997); Forester et al., *Exp. Agric.* **33**:15-33 (1997). Minor variations make these technologies applicable to a broad range of plant species.

Each of these techniques has advantages and disadvantages. In each of the techniques, DNA from a plasmid is genetically engineered such that it contains not only the gene of interest, but also selectable and screenable marker genes. A selectable marker gene is used to select only those cells that have integrated copies of the plasmid (the construction is such that the gene of interest and the selectable and screenable genes are transferred as a unit). The screenable gene provides another check for the successful culturing of only those cells carrying the genes of interest. A commonly used selectable marker gene is neomycin phosphotransferase II (NPT II). This gene conveys resistance to kanamycin, a compound that can be added directly to the growth media on which the cells grow. Plant cells are normally susceptible to kanamycin and, as a result, die. The presence of the NPT II gene overcomes the effects of the kanamycin and each cell with this gene remains viable. Another selectable marker gene which can be employed in the practice of this invention is the gene which confers resistance to the herbicide glufosinate (Basta). A screenable gene commonly used is the β -glucuronidase gene (GUS). The presence of this gene is characterized using a histochemical reaction in which a sample of putatively transformed cells is treated with a GUS assay solution. After an appropriate incubation, the cells containing the GUS gene turn blue.

The plasmid containing one or more of these genes is introduced into either plant protoplasts or callus cells by any of the previously mentioned techniques. If the marker gene is a selectable gene, only those cells that have incorporated the DNA package survive under selection with the appropriate phytotoxic agent. Once the

appropriate cells are identified and propagated, plants are regenerated. Progeny from the transformed plants must be tested to insure that the DNA package has been successfully integrated into the plant genome.

Mammalian host cells may also be used in the practice of the invention.

- 5 Examples of suitable mammalian cell lines include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line 293S (Graham et al., *J. Gen. Virol.* **36**:59 [1977]); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells (Urlab and Chasin, *Proc. Natl. Acad. Sci. USA* **77**:4216 [1980]); mouse sertoli cells (TM4, Mather, *Biol. Reprod.* **23**:243 [1980]); monkey kidney cells (CV1-76, ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse
- 10 mammary tumor cells (MMT 060562, ATCC CCL 51); rat hepatoma cells (HTC, ML54, Baumann et al., *J. Cell Biol.* **85**:1 [1980]); and TRI cells (Mather et al., *Annals N.Y. Acad. Sci.* **383**:44 [1982]). Expression vectors for these cells ordinarily include (if necessary) DNA sequences for an origin of replication, a promoter located in front of the gene to be expressed, a ribosome binding site, an RNA splice site, a
- 15 polyadenylation site, and a transcription terminator site.
- 20

Promoters used in mammalian expression vectors are often of viral origin. These viral promoters are commonly derived from polyoma virus, Adenovirus 2, and most frequently Simian Virus 40 (SV40). The SV40 virus contains two promoters that are termed the early and late promoters. These promoters are particularly useful

25 because they are both easily obtained from the virus as one DNA fragment that also contains the viral origin of replication (Fiers et al., *Nature* **273**:113 [1978]). Smaller or larger SV40 DNA fragments may also be used, provided they contain the approximately 250-bp sequence extending from the HindIII site toward the BglI site located in the viral origin of replication.

30 Alternatively, promoters that are naturally associated with the foreign gene (homologous promoters) may be used provided that they are compatible with the host cell line selected for transformation.

An origin of replication may be obtained from an exogenous source, such as SV40 or other virus (e.g., Polyoma, Adeno, VSV, BPV) and inserted into the cloning

35 vector. Alternatively, the origin of replication may be provided by the host cell

chromosomal replication mechanism. If the vector containing the foreign gene is integrated into the host cell chromosome, the latter is often sufficient.

5 The use of a secondary DNA coding sequence can enhance production levels of (-)-limonene synthase, (-)-pinene synthase and myrcene synthase in transformed cell lines. The secondary coding sequence typically comprises the enzyme dihydrofolate reductase (DHFR). The wild-type form of DHFR is normally inhibited by the chemical methotrexate (MTX). The level of DHFR expression in a cell will vary depending on the amount of MTX added to the cultured host cells. An additional feature of DHFR that makes it particularly useful as a secondary sequence 10 is that it can be used as a selection marker to identify transformed cells. Two forms of DHFR are available for use as secondary sequences, wild-type DHFR and MTX-resistant DHFR. The type of DHFR used in a particular host cell depends on whether the host cell is DHFR deficient (such that it either produces very low levels of DHFR endogenously, or it does not produce functional DHFR at all). DHFR-deficient cell 15 lines such as the CHO cell line described by Urlaub and Chasin, *supra*, are transformed with wild-type DHFR coding sequences. After transformation, these DHFR-deficient cell lines express functional DHFR and are capable of growing in a culture medium lacking the nutrients hypoxanthine, glycine and thymidine. Nontransformed cells will not survive in this medium.

20 The MTX-resistant form of DHFR can be used as a means of selecting for transformed host cells in those host cells that endogenously produce normal amounts of functional DHFR that is MTX sensitive. The CHO-K1 cell line (ATCC No. CL 61) possesses these characteristics, and is thus a useful cell line for this purpose. The addition of MTX to the cell culture medium will permit only those 25 cells transformed with the DNA encoding the MTX-resistant DHFR to grow. The nontransformed cells will be unable to survive in this medium.

Prokaryotes may also be used as host cells for the initial cloning steps of this invention. They are particularly useful for rapid production of large amounts of DNA, for production of single-stranded DNA templates used for site-directed 30 mutagenesis, for screening many mutants simultaneously, and for DNA sequencing of the mutants generated. Suitable prokaryotic host cells include *E. coli* K12 strain 94 (ATCC No. 31,446), *E. coli* strain W3110 (ATCC No. 27,325) *E. coli* X1776 (ATCC No. 31,537), and *E. coli* B; however many other strains of *E. coli*, such as HB101, JM101, NM522, NM538, NM539, and many other species and 35 genera of prokaryotes including bacilli such as *Bacillus subtilis*, other

enterobacteriaceae such as *Salmonella typhimurium* or *Serratia marcesans*, and various *Pseudomonas* species may all be used as hosts. Prokaryotic host cells or other host cells with rigid cell walls are preferably transformed using the calcium chloride method as described in section 1.82 of Sambrook et al., *supra*.
5 Alternatively, electroporation may be used for transformation of these cells. Prokaryote transformation techniques are set forth in Dower, W.J., in Genetic Engineering, Principles and Methods, 12:275-296, Plenum Publishing Corp., 1990; Hanahan et al., *Meth. Enzymol.*, 204:63 (1991).

As a representative example, cDNA sequences encoding (-)-limonene
10 synthase, (-)-pinene synthase or myrcene synthase may be transferred to the (His)₆-Tag pET vector commercially available (from Novagen) for overexpression in *E. coli* as heterologous host. This pET expression plasmid has several advantages in high level heterologous expression systems. The desired cDNA insert is ligated in frame to plasmid vector sequences encoding six histidines followed by a highly
15 specific protease recognition site (thrombin) that are joined to the amino terminus codon of the target protein. The histidine "block" of the expressed fusion protein promotes very tight binding to immobilized metal ions and permits rapid purification of the recombinant protein by immobilized metal ion affinity chromatography. The histidine leader sequence is then cleaved at the specific proteolysis site by treatment
20 of the purified protein with thrombin, and the (-)-limonene synthase, (-)-pinene synthase and myrcene synthase again purified by immobilized metal ion affinity chromatography, this time using a shallower imidazole gradient to elute the recombinant synthases while leaving the histidine block still adsorbed. This overexpression-purification system has high capacity, excellent resolving power and
25 is fast, and the chance of a contaminating *E. coli* protein exhibiting similar binding behavior (before and after thrombin proteolysis) is extremely small.

As will be apparent to those skilled in the art, any plasmid vectors containing replicon and control sequences that are derived from species compatible with the host cell may also be used in the practice of the invention. The vector usually has a
30 replication site, marker genes that provide phenotypic selection in transformed cells, one or more promoters, and a polylinker region containing several restriction sites for insertion of foreign DNA. Plasmids typically used for transformation of *E. coli* include pBR322, pUC18, pUC19, pUC118, pUC119, and Bluescript M13, all of which are described in sections 1.12-1.20 of Sambrook et al., *supra*. However, many
35 other suitable vectors are available as well. These vectors contain genes coding for

ampicillin and/or tetracycline resistance which enables cells transformed with these vectors to grow in the presence of these antibiotics.

The promoters most commonly used in prokaryotic vectors include the β -lactamase (penicillinase) and lactose promoter systems (Chang et al., *Nature* **375**:615 [1978]; Itakura et al., *Science* **198**:1056 [1977]; Goeddel et al., *Nature* **281**:544 [1979]) and a tryptophan (trp) promoter system (Goeddel et al., *Nucl. Acids Res.* **8**:4057 [1980]; EPO Appl. Publ. No. 36,776), and the alkaline phosphatase systems. While these are the most commonly used, other microbial promoters have been utilized, and details concerning their nucleotide sequences have been published, enabling a skilled worker to ligate them functionally into plasmid vectors (see Siebenlist et al., *Cell* **20**:269 [1980]).

Many eukaryotic proteins normally secreted from the cell contain an endogenous secretion signal sequence as part of the amino acid sequence. Thus, proteins normally found in the cytoplasm can be targeted for secretion by linking a signal sequence to the protein. This is readily accomplished by ligating DNA encoding a signal sequence to the 5' end of the DNA encoding the protein and then expressing this fusion protein in an appropriate host cell. The DNA encoding the signal sequence may be obtained as a restriction fragment from any gene encoding a protein with a signal sequence. Thus, prokaryotic, yeast, and eukaryotic signal sequences may be used herein, depending on the type of host cell utilized to practice the invention. The DNA and amino acid sequence encoding the signal sequence portion of several eukaryotic genes including, for example, human growth hormone, proinsulin, and proalbumin are known (see Stryer, *Biochemistry* W.H. Freeman and Company, New York, NY, p. 769 [1988]), and can be used as signal sequences in appropriate eukaryotic host cells. Yeast signal sequences, as for example acid phosphatase (Arima et al., *Nucl. Acids Res.* **11**:1657 [1983]), α -factor, alkaline phosphatase and invertase may be used to direct secretion from yeast host cells. Prokaryotic signal sequences from genes encoding, for example, LamB or OmpF (Wong et al., *Gene* **68**:193 [1988]), MalE, PhoA, or beta-lactamase, as well as other genes, may be used to target proteins from prokaryotic cells into the culture medium.

Trafficking sequences from plants, animals and microbes can be employed in the practice of the invention to direct the monoterpene synthase proteins of the present invention to the cytoplasm, endoplasmic reticulum, mitochondria or other cellular components, or to target the protein for export to the medium. These considerations apply to the overexpression of (-)-limonene synthase, (-)-pinene

synthase and myrcene synthase, and to direction of expression within cells or intact organisms to permit gene product function in any desired location.

5 The construction of suitable vectors containing DNA encoding replication sequences, regulatory sequences, phenotypic selection genes and the monoterpene synthase DNA of interest are prepared using standard recombinant DNA procedures. Isolated plasmids and DNA fragments are cleaved, tailored, and ligated together in a specific order to generate the desired vectors, as is well known in the art (see, for example, Maniatis, *supra*, and Sambrook et al., *supra*).

10 As discussed above, (-)-limonene synthase, (-)-pinene synthase and myrcene synthase variants are preferably produced by means of mutation(s) that are generated using the method of site-specific mutagenesis. This method requires the synthesis and use of specific oligonucleotides that encode both the sequence of the desired mutation and a sufficient number of adjacent nucleotides to allow the oligonucleotide to stably hybridize to the DNA template.

15 The foregoing may be more fully understood in connection with the following representative examples, in which "Plasmids" are designated by a lower case p followed by an alphanumeric designation. The starting plasmids used in this invention are either commercially available, publicly available on an unrestricted basis, or can be constructed from such available plasmids using published procedures. In addition, other equivalent plasmids are known in the art and will be apparent to the ordinary artisan.

20 "Digestion", "cutting" or "cleaving" of DNA refers to catalytic cleavage of the DNA with an enzyme that acts only at particular locations in the DNA. These enzymes are called restriction endonucleases, and the site along the DNA sequence where each enzyme cleaves is called a restriction site. The restriction enzymes used in this invention are commercially available and are used according to the instructions supplied by the manufacturers. (See also sections 1.60-1.61 and sections 3.38-3.39 of Sambrook et al., *supra*.)

25 "Recovery" or "isolation" of a given fragment of DNA from a restriction digest means separation of the resulting DNA fragment on a polyacrylamide or an agarose gel by electrophoresis, identification of the fragment of interest by comparison of its mobility versus that of marker DNA fragments of known molecular weight, removal of the gel section containing the desired fragment, and separation of the gel from DNA. This procedure is known generally. For example, see Lawn et al.

(*Nucleic Acids Res.* **9**:6103-6114 [1982]), and Goeddel et al. (*Nucleic Acids Res.*, *supra*).

The following examples merely illustrate the best mode now contemplated for practicing the invention, but should not be construed to limit the invention. All literature citations herein are expressly incorporated by reference.

EXAMPLE 1

PCR-Based Generation of Probes for Cloning Monoterpene Synthases From Grand fir (*Abies grandis*)

Substrates, Reagents and cDNA Library – [$1\text{-}^3\text{H}$]Geranyl diphosphate (250 Ci/mol) (Croteau, R., Alonso, W.R., Koepp, A.E., and Johnson, M.A. (1994) *Arch. Biochem. Biophys.* **309**:184-192), [$1\text{-}^3\text{H}$]farnesyl diphosphate (125 Ci/mol) (Dehal, S.S., and Croteau, R. (1988) *Arch. Biochem. Biophys.* **261**:346-356) and [$1\text{-}^3\text{H}$]geranylgeranyl diphosphate (120 Ci/mol) (LaFever, R.E., Stofer Vogel, B., and Croteau, R. (1994) *Arch. Biochem. Biophys.* **313**:139-149) were prepared as described previously. Terpenoid standards were from our own collection. All other biochemicals and reagents were purchased from Sigma Chemical Co. or Aldrich Chemical Co., unless otherwise noted. Construction of the λ ZAP II cDNA library, using mRNA isolated from wounded Grand fir sapling stems, was described previously (Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268).

PCR-Based Probe Generation – Based on comparison of sequences of limonene synthase from spearmint (Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* **268**:23016-23024), 5-*epi*-aristolochene synthase from tobacco (Facchini, P.J., and Chappell, J. (1992) *Proc. Natl. Acad. Sci. USA* **89**:11088-11092), and casbene synthase from castor bean (Mau, C.J.D., and West, C.A. (1994) *Proc. Natl. Acad. Sci. USA* **91**:8497-8501), four conserved regions were identified for which a set of consensus degenerate primers were synthesized: Primer A (SEQ ID NO:7); Primer B (SEQ ID NO:8); Primer C (SEQ ID NO:9); Primer D (SEQ ID NO:10). Primers A (SEQ ID NO:7), B (SEQ ID NO:8) and C (SEQ ID NO:9) have been described previously (Steele, C., Lewinsohn, E. and Croteau, R., *Proc. Nat'l. Acad. Sci. USA*, **92**: 4164-4168 (1995)); primer D (SEQ ID NO:10) was designed based on the conserved amino acid sequence motif DD(T/I)(I/Y/F)D(A/V)Y(A/G)(SEQ ID NO:25) of the above noted terpene synthases (Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* **268**:23016-23024;

Facchini, P.J., and Chappell, J. (1992) *Proc. Natl. Acad. Sci. USA* **89**:11088-11092; Mau, C.J.D., and West, C.A. (1994) *Proc. Natl. Acad. Sci. USA* **91**:8497-8501).

Each of the sense primers, A (SEQ ID NO:7), B (SEQ ID NO:8) and D (SEQ ID NO:10), was used for PCR in combination with antisense primer C (SEQ ID NO:9) by employing a broad range of amplification conditions. PCR was performed in a total volume of 50 μ l containing 20 mM Tris/HCl (tris(hydroxymethyl) aminomethane/HCl, pH 8.4), 50 mM KCl, 5 mM MgCl₂, 200 μ M of each dNTP, 1-5 μ M of each primer, 2.5 units of *Taq* polymerase (BRL) and 5 μ l of purified Grand fir stem cDNA library phage as template (1.5×10^9 pfu/ml). Analysis of the PCR reaction products by agarose gel electrophoresis (Sambrock, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY) revealed that only the combination of primers C and D generated a specific PCR product of approximately 110 bps (base pairs). This PCR product was gel purified, ligated into pT7Blue (Novagen), and transformed into *E. coli* XL1-Blue cells. Plasmid DNA was prepared from 41 individual transformants and the inserts were sequenced (DyeDeoxy Terminator Cycle Sequencing, Applied Biosystems). Four different insert sequences were identified, and were designated as probes 1 (SEQ ID NO:11), 2 (SEQ ID NO:12), 4 (SEQ ID NO:13) and 5 (SEQ ID NO:14).

Subsequent isolation of four new cDNA species (*AG1.28* (SEQ ID NO:15); *AG2.2* (SEQ ID NO:16); *AG4.30* (SEQ ID NO:17) and *AG5.9* (SEQ ID NO:19)), encoding terpene synthases from Grand fir corresponding to probes 1 (SEQ ID NO:11), 2 (SEQ ID NO:12), 4 (SEQ ID NO:13) and 5 (SEQ ID NO:14), respectively, allowed the identification of three additional conserved sequence elements which were used to design a set of three new PCR primers: Primer E (5'-GGI GA(A/G) A(A/C)(A/G) (A/G)TI ATG GA(A/G) GA(A/G) GC-3')(SEQ ID NO:21); Primer F (5'-GA(A/G) (C/T)TI CA(G/A) (C/T)TI (A/C/T)(C/G/T)I (A/C)GI TGG TGG-3')(SEQ ID NO:22) and Primer G (5'-CCA (A/G)TT IA(A/G) ICC (C/T)TT IAC (A/G)TC-3')(SEQ ID NO:23).

Degenerate primer E (SEQ ID NO:21) was designed to conserved element GE(K/T)(V/I)M(E/D)EA (SEQ ID NO:26) and degenerate primer F (SEQ ID NO:22) was designed to conserved element Q(F/Y/D)(I/L)(T/L/R)RWW (SEQ ID NO:27) by comparing the sequences of five cloned terpene synthases from Grand fir: a monoterpene synthase corresponding to probe 2 (SEQ ID NO: 12), two sesquiterpene synthases corresponding to probe 4 (SEQ ID NO:13) and probe 5 (SEQ ID NO:14),

respectively, a previously described diterpene synthase (Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268), and a truncated terpene synthase corresponding to probe 1 (SEQ ID NO:11). Degenerate primer G (SEQ ID NO:23) was designed according to the amino acid sequence DVIKG(F/L)NW (SEQ ID NO:28) obtained from a peptide generated by trypsin digestion of purified (-)-pinene synthase from Grand fir. Primers E (SEQ ID NO:21) and F (SEQ ID NO:22) were independently used for PCR amplification in combination with primer G (SEQ ID NO:23), with Grand fir stem cDNA library as template. The combination of primers E (SEQ ID NO:21) and G (SEQ ID NO:23) yielded a specific PCR product of approximately 1020 bps. This PCR product was ligated into pT7Blue and transformed into *E. coli* XL1-Blue. Plasmid DNA was prepared from 20 individual transformants and inserts were sequenced from both ends. The sequence of this 1022 bp insert was identical for all 20 plasmids and was designated as probe 3 (SEQ ID NO:24).

EXAMPLE 2

Screening a Wounded Grand fir Stem cDNA Library

For library screening, 100 ng of each probe was amplified by PCR, gel purified, randomly labeled with [α -³²P]dATP (Feinberg, A.P., and Vogelstein, B. (1984) *Anal. Biochem.* **137**:266-267), and used individually to screen replica filters of 10⁵ plaques of the wound-induced Grand fir stem cDNA library plated on *E. coli* LE392. Hybridization with probes 1 (SEQ ID NO:11), 2 (SEQ ID NO:12), 4 (SEQ ID NO:13) and 5 (SEQ ID NO:14) was performed for 14 h at 65 °C in 3 x SSPE and 0.1% SDS. Filters were washed three times for 10 min at 55 °C in 3 x SSPE with 0.1% SDS and exposed for 12 h to Kodak XAR film at -70 °C. All of the λ ZAPII clones yielding positive signals were purified through a second round of hybridization (probe 1 (SEQ ID NO:11) gave 25 positives, probe 2 (SEQ ID NO:12) gave 16 positives, probe 4 (SEQ ID NO:13) gave 49 positives and probe 5 (SEQ ID NO:14) gave 12 positives).

Hybridization with probe 3 (SEQ ID NO:24) was performed as before, but the filters were washed three times for 10 min at 65 °C in 3 x SSPE and 0.1% SDS before exposure. Approximately 400 λ ZAPII clones yielded strong positive signals, and 34 of these were purified through a second round of hybridization at 65°C. Approximately 400 additional clones yielded weak positive signals with probe 3 (SEQ ID NO:24), and 18 of these were purified through a second round of hybridization for 20 h at 45 °C. Purified λ ZAP II clones isolated using all five

probes were *in vivo* excised as Bluescript II SK(-) phagemids and transformed into *E. coli* XL0LR according to the manufacturer's instructions (Stratagene). The size of each cDNA insert was determined by PCR using T3 (SEQ ID NO:29) and T7 (SEQ ID NO:30) promoter primers and selected inserts (>1.5 kb) were partially sequenced from both ends.

EXAMPLE 3

Grand fir Monoterpene Synthase cDNA Expression in *E. coli* and Enzyme

Assays

Except for cDNA clones AG3.18 (SEQ ID NO:3) and AG3.48 (SEQ ID NO:31), all of the partially sequenced inserts were either truncated at the 5'-end, or were out of frame, or bore premature stop codons upstream of the presumptive methionine start codon. For the purpose of functional expression, a 2023 bp insert fragment, extending from nucleotides 75 to 2097 of the sequence set forth in SEQ ID NO:1, and a 1911 bp insert fragment, extending from nucleotide 1 to nucleotide 1910 of the sequence set forth in SEQ ID NO:3, were subcloned in frame into pGEX vectors (Pharmacia). A 2016 bp fragment extending from nucleotide 73 to nucleotide 2088 of the sequence set forth in SEQ ID NO:5 was subcloned in frame into the pSBETa vector (Schenk, P.M., Baumann, S., Mattes, R., and Steinbiss, H.-H. (1995) *Biotechniques* 19, 196-200). To introduce suitable restriction sites for subcloning, fragments were amplified by PCR using primer combinations 2.2-*Bam*HI (5'-CAA AGG GAT CCA GAA TGG CTC TGG-3')(SEQ ID NO:33) and 2.2-*Not*I (5'-AGT AAG CGG CCG CTT TTT AAT CAT ACC CAC-3')(SEQ ID NO:34) with pAG2.2 insert (SEQ ID NO:1) as template, 3.18-*Eco*RI (5'-CTG CAG GAA TTC GGC ACG AGC-3')(SEQ ID NO:35) and 3.18-*Sma*I (5'-CAT AGC CCC GGG CAT AGA TTT GAG CTG-3')(SEQ ID NO:36) with pAG3.18 insert (SEQ ID NO:3) as template, and 10-*Nde*I (5'-GGC AGG AAC ATA TGG CTC TCC TTT CTA TCG-3')(SEQ ID NO:37) and 10-*Bam*HI (5'-TCT AGA ACT AGT GGATCC CCC GGC CTG CAG-3')(SEQ ID NO:38) with pAG10 insert (SEQ ID NO:5) as template.

PCR reactions were performed in volumes of 50 μ l containing 20 mM Tris/HCl (pH 8.8), 10 mM KCl, 10 mM $(\text{NH}_4)_2\text{SO}_4$, 2 mM MgSO_4 , 0.1% Triton X-100, 5 μ g bovine serum albumin (BSA), 200 μ M of each dNTP, 0.1 μ M of each primer, 2.5 units of recombinant *Pfu* polymerase (Stratagene) and 100 ng plasmid DNA with the following program: denaturation at 94 $^\circ\text{C}$, 1 min; annealing at 60 $^\circ\text{C}$, 1 min; extension at 72 $^\circ\text{C}$, 3.5 min; 35 cycles with final extension at 72 $^\circ\text{C}$, 5 min. The PCR products were purified by agarose gel electrophoresis and used as

template for a secondary PCR amplification with the identical conditions in total volumes of 250 μ l each. Products from this secondary amplification were digested with the above indicated restriction enzymes, purified by ultrafiltration and then ligated, respectively, into *Bam*HI/*Not*I-digested pGEX-4T-2 to yield plasmid pGAG2.2, into *Eco*RI/*Sma*I-digested pGEX-4T-3 to yield plasmid pGAG3.18, and into *Nde*I/*Bam*HI-digested pSBETa to yield plasmid pSBAG10; these plasmids were then transformed into *E. coli* XL1-Blue or *E. coli* BL21(DE3).

For expression, bacterial strains *E. coli* XL0LR/pAG3.18, *E. coli* XL0LR/pAG3.48, *E. coli* XL1-Blue/pGAG2.2, *E. coli* XL1-Blue/pGAG3.18, and *E. coli* BL21(DE3)/pSBAG10 were grown to $A_{600} = 0.5$ at 37 °C in 5 ml of LB medium (Sambrock, J., Fritsch, E. F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY) supplemented with 100 μ g ampicillin/ml or 30 μ g kanamycin/ml as determined by the vector. Cultures were then induced by addition of 1 mM isopropyl-1-thio- β -D-galactopyranoside and grown for another 12 h at 20 °C. Cells were harvested by centrifugation (2000 x g, 10 min) and resuspended in either 1 ml monoterpene synthase assay buffer [50 mM Tris/HCl (pH 7.5), 500 mM KCl, 1 mM $MnCl_2$, 5 mM dithiothreitol, 0.05% (w/v) $NaHSO_3$ and 10% (v/v) glycerol], 1 ml sesquiterpene synthase assay buffer [10 mM dibasic potassium phosphate, 1.8 mM monobasic potassium phosphate (pH 7.3), 140 mM NaCl, 10 mM $MgCl_2$, 5 mM dithiothreitol, 0.05% (w/v) $NaHSO_3$ and 10% (v/v) glycerol], or 1 ml diterpene synthase assay buffer [30 mM Hepes (*N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid, pH 7.2), 7.5 mM $MgCl_2$, 5 mM dithiothreitol, 10 μ M $MnCl_2$, 0.05% (w/v) $NaHSO_3$ and 10% (v/v) glycerol].

Cells were disrupted by sonication (Braun-Sonic 2000 with microprobe at maximum power for 15 seconds at 0-4°C), the homogenates were cleared by centrifugation (18,000 x g, 10 min), and 1 ml of the resulting supernatant was assayed for monoterpene synthase activity with 2.5 μ M of [$1-^3H$]geranyl diphosphate, for sesquiterpene synthase activity with 3.5 μ M [$1-^3H$]farnesyl diphosphate, or for diterpene synthase activity with 5 μ M [$1-^3H$]geranylgeranyl diphosphate following standard protocols (Croteau, R., and Cane, D.E. (1985) *Methods Enzymol.* **110**:383-405; LaFever, R.E., Stofer Vogel, B., and Croteau, R. (1994) *Arch. Biochem. Biophys.* **313**:139-149; Dehal, S.S., and Croteau, R. (1988) *Arch. Biochem. Biophys.* **261**:346-356). In the case of the monoterpene synthase and sesquiterpene synthase assays, the incubation mixture was overlaid with 1 ml pentane

to trap volatile products. In all cases, after incubation at 31°C for 2 h, the reaction mixture was extracted with pentane (3 x 1 ml) and the combined extract was passed through a 1.5 ml column of anhydrous MgSO₄ and silica gel (Mallinckrodt 60 Å) to provide the terpene hydrocarbon fraction free of oxygenated metabolites. The columns were subsequently eluted with 3 x 1 ml of ether to collect any oxygenated products, and an aliquot of each fraction was taken for liquid scintillation counting to determine conversion rate.

Product Identification – To obtain sufficient product for analysis by radio-GLC (gas liquid chromatography), chiral capillary GLC and GLC-MS (mass spectrum/spectrometry), preparative-scale enzyme incubations were carried out. Thus, the enzyme was prepared from 50 ml of cultured bacterial cells by extraction with 3 ml of assay buffer as above, and the extracts were incubated with excess substrate overnight at 31°C. The hydrocarbon fraction was isolated by elution through MgSO₄-silica gel as before, and the pentane eluate was concentrated for evaluation by capillary radio-GLC as described (Croteau, R., and Satterwhite, D.M. (1990) *J. Chromatogr.* **500**:349-354), by chiral column capillary GLC (Lewinsohn, E., Savage, T.J., Gijzen, M., and Croteau, R. (1993) *Phytochem. Anal.* **4**:220-225), and by combined GLC-MS [Hewlett-Packard 6890 GC-MSD with cool (40°C) on-column injection, detection via electron impact ionization (70 eV), He carrier at 0.7 psi., column: 0.25 mm i.d. x 30 m fused silica with 0.25 µm film of 5MS (Hewlett-Packard) programmed from 35°C (5 min hold) to 230°C at 5°C/min].

EXAMPLE 4

Sequence Analysis

Inserts of all recombinant bluescript plasmids and pGEX plasmids were completely sequenced on both strands via primer walking and nested deletions (Sambrock, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY) using the DyeDeoxy Terminator Cycle Sequencing method (Applied Biosystems). Sequence analysis was done using the Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, WI.

EXAMPLE 5

RNA Extraction and Northern Blotting

Grand fir sapling stem tissue was harvested prior to wounding or two days after wounding by a standard procedure (Gijzen, M., Lewinsohn, E., and Croteau, R. (1991) *Arch. Biochem. Biophys.* **289**:267-273). Total RNA was isolated (Lewinsohn,

E., Steele, C.L., and Croteau, R. (1994) *Plant Mol. Biol. Rep.* **12**:20-25) and 20 µg of RNA per gel lane was separated under denaturing conditions (Sambrock, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY) and transferred to nitrocellulose membranes (Schleicher and Schuell) according to the manufacturer's protocol. To prepare hybridization probes, cDNA fragments of 1.4-1.5 kb were amplified by PCR from AG2.2 (SEQ ID NO:1) with primer JB29 (5'-CTA CCA TTC CAA TAT CTG-3')(SEQ ID NO:39) and primer 2-8 (5'-GTT GGA TCT TAG AAG TTC CC-3')(SEQ ID NO:40), from AG3.18 (SEQ ID NO:3) with primer 3-9 (5'-TTT CCA TTC CAA CCT CTG GG-3')(SEQ ID NO:41) and primer 3-11 (5'-CGT AAT GGA AAG CTC TGG CG-3')(SEQ ID NO:42), and from AG10 (SEQ ID NO:5) with primer 7-1 (5'-CCT TAC ACG CCT TTG GAT GG-3')(SEQ ID NO:43) and primer 7-3 (5'-TCT GTT GAT CCA GGA TGG TC-3')(SEQ ID NO:44). The probes were randomly labeled with [α -³²P]dATP (Feinberg, A.P., and Vogelstein, B. (1984) *Anal. Biochem.* **137**:266-267). Blots were hybridized for 24 h at 55°C in 3 x SSPE and 0.1% SDS, washed at 55°C in 1 x SSPE and 0.1% SDS and subjected to autoradiography as described above at -80°C for 24 h.

EXAMPLE 6

Cloning and Characterization of Clones AG1.28 (SEQ ID NO:15), AG2.2 (SEQ ID NO:1), AG4.30 (SEQ ID NO:17) and AG 5.9 (SEQ ID NO:19)

Similarity-Based Cloning of Grand fir Terpene Synthases – Grand fir has been developed as a model system for the study of induced oleoresin production in conifers in response to wounding and insect attack (Johnson, M.A., and Croteau, R. (1987) in *Ecology and Metabolism of Plant Lipids* (Fuller, G., and Nes, W.D., eds) pp. 76-91, American Chemical Society Symposium Series 325, Washington, DC; Gijzen, M., Lewinsohn, E., Savage, T.J., and Croteau, R.B. (1993) in *Bioactive Volatile Compounds from Plants* (Teranishi, R., Buttery, R.G., and Sugisawa, H., eds) pp. 8-22, American Chemical Society Symposium Series 525, Washington, DC; Raffa, K. F., and Berryman, A.A. (1982) *Can. Entomol.* **114**:797-810; Steele, C., Lewinsohn, E., and Croteau, R. (1995) *Proc. Natl. Acad. Sci. USA* **92**:4164-4168; Lewinsohn, E., Gijzen, M., and Croteau, R. (1992) in *Regulation of Isopentenoid Metabolism* (Nes, W.D., Parish, E.J., and Trzaskos, J.M., eds) pp. 8-17, American Chemical Society Symposium Series 497, Washington, DC). The chemistry and biosynthesis of the oleoresin monoterpenes, sesquiterpenes and diterpenes have been well defined (Lewinsohn, E., Savage, T.J., Gijzen, M., and

- Croteau, R. (1993) *Phytochem. Anal.* **4**:220-225; Lewinsohn, E., Gijzen, M., and Croteau, R. (1991) *Plant Physiol.* **96**:44-49; Funk, C., Lewinsohn, E., Stofer Vogel, B., Steele C., and Croteau, R. (1994) *Plant Physiol.* **106**:999-1005; Gijzen, M., Lewinsohn, E., and Croteau, R. (1991) *Arch. Biochem. Biophys.* **289**:267-273;
- 5 Lewinsohn, E., Gijzen, M., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **293**:167-173; LaFever, R.E., Stofer Vogel, B., and Croteau, R. (1994) *Arch. Biochem. Biophys.* **313**:139-149; Funk, C., and Croteau, R. (1994) *Arch. Biochem. Biophys.* **308**:258-266); however, structural analysis of the responsible terpene synthases as well as studies on the regulation of oleoresinosis require the isolation of
- 10 cDNA species encoding the terpene synthases. Protein purification from conifers, as the basis for cDNA isolation, has been of limited success (Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268) and thus far has not permitted cloning of any of the monoterpene synthases from these species (Steele, C., Lewinsohn, E., and Croteau, R. (1995) *Proc. Natl. Acad. Sci. USA* **92**:4164-4168).
- 15

As a possible alternative to protein-based cloning of terpene synthases, a homology-based PCR strategy was proposed (Steele, C., Lewinsohn, E., and Croteau, R. (1995) *Proc. Natl. Acad. Sci. USA* **92**:4164-4168) that was founded upon the three terpene synthases of plant origin then available, a monoterpene synthase,

20 (-)-4S-limonene synthase, from spearmint (*Mentha spicata*, Lamiaceae) (Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* **268**:23016-23024), a sesquiterpene synthase, 5-*epi*-aristolochene synthase, from tobacco (*Nicotiana tabacum*, Solanaceae) (Facchini, P.J., and Chappell, J. (1992) *Proc. Natl. Acad. Sci. USA* **89**:11088-11092), and a diterpene synthase,

25 casbene synthase, from castor bean (*Ricinus communis*, Euphorbiaceae) (Mau, C.J.D., and West, C.A. (1994) *Proc. Natl. Acad. Sci. USA* **91**:8497-8501). Despite the taxonomic distances between these three angiosperm species and the differences in substrate utilized, reaction mechanism and product type of the three enzymes, a comparison of the deduced amino acid sequences identified several conserved

30 regions that appeared to be useful for the design of degenerate PCR primers (see Example 1). Using cDNA from a wound-induced Grand fir stem library as template, PCR primers C (SEQ ID NO:9) and D (SEQ ID NO:10) amplified products corresponding to four distinct sequence groups, all of which showed significant similarity to sequences of cloned terpene synthases of plant origin. The four

35 different inserts were designated as probes 1 (SEQ ID NO:11), 2 (SEQ ID NO:12), 4

(SEQ ID NO:13) and 5 (SEQ ID NO:14), and were employed for isolation of the corresponding cDNA clones by plaque hybridization.

Screening of 10^5 cDNA phage plaques from the wounded Grand fir stem library, with each of the four probes, yielded a four-fold difference in the number of positives, most likely reflecting different levels of expression of the corresponding genes. Size selected inserts (>1.5 kb) of purified and *in vivo* excised clones were partially sequenced from both ends, and were shown to segregate into four distinct groups corresponding to the four hybridization probes. Since all cDNAs corresponding to probes 1 (SEQ ID NO:11), 4 (SEQ ID NO:13) and 5 (SEQ ID NO:14) were truncated at their 5'-ends, only inserts of the largest representatives of each group, clone *AG1.28* (SEQ ID NO:15), clone *AG2.2* (SEQ ID NO:1) (apparently full length), clone *AG4.30* (SEQ ID NO:17) and clone *AG5.9* (SEQ ID NO:19), were completely sequenced. Clone *AG1.28* (SEQ ID NO:15)(2424 bps) includes an open reading frame (ORF) of 2350 nucleotides (nts) encoding 782 amino acids (SEQ ID NO:16); clone *AG2.2* (SEQ ID NO:1)(2196 bps), includes an ORF of 1881 nts encoding 627 amino acids (SEQ ID NO:2); clone *AG4.30* (SEQ ID NO:17)(1967 bps) includes an ORF of 1731 nts encoding 577 amino acids (SEQ ID NO:18) and clone *AG5.9* (SEQ ID NO:19)(1416 bps) includes an ORF of 1194 nucleotides encoding 398 amino acids (SEQ ID NO:20).

cDNA clones *AG1.28* (SEQ ID NO:15), *AG2.2* (SEQ ID NO:1), *AG4.30* (SEQ ID NO:17) and *AG5.9* (SEQ ID NO:19) were compared pairwise with each other and with other cloned plant terpene synthases. Truncated clone *AG1.28* (SEQ ID NO:15) resembled most closely in size and sequence (72% similarity, 49% identity) a diterpene cyclase, abietadiene synthase, from Grand fir (Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268). Clones *AG4.30* (SEQ ID NO:17) and *AG5.9* (SEQ ID NO:19) share approximately 80% similarity (60% identity) at the amino acid level, and are almost equally distant from both clone *AG1.28* (SEQ ID NO:15) and full-length clone *AG2.2* (SEQ ID NO:1)(range of 65-70% similarity and 45-47% identity); the amino acid sequence similarity between *AG1.28* (SEQ ID NO:15) and *AG2.2* (SEQ ID NO:1) is 65% (41% identity). Considering the high level of homology between *AG4.30* (SEQ ID NO:17) and *AG5.9* (SEQ ID NO:19), these comparisons suggest that the four new cDNAs, *AG1.28* (SEQ ID NO:15), *AG2.2* (SEQ ID NO:1), *AG4.30* (SEQ ID NO:17) and *AG5.9* (SEQ ID NO:19), represent the three major subfamilies

of Grand fir terpene synthase genes encoding monoterpene synthases, sesquiterpene synthases and diterpene synthases.

Identification of cDNA Clone AG2.2 (SEQ ID NO:1) as Myrcene Synthase

- The pAG2.2 insert (SEQ ID NO:1) appeared to be a full-length clone encoding a protein of molecular weight 72,478 with a calculated pI at 6.5. The size of the translated protein encoded by AG2.2 (SEQ ID NO:1) (627 residues)(SEQ ID NO:2) is in the range of the monoterpene synthase preproteins for limonene synthase from spearmint (Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* **268**:23016-23024) and *Perilla frutescens* (Yuba, A., Yazaki, K., Tabata, M., Honda, G., and Croteau, R. (1996) *Arch. Biochem. Biophys.* **332**:280-287), but is about 240 amino acids shorter than the two gymnosperm diterpene synthase preproteins for abietadiene synthase (Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268) and taxadiene synthase (Wildung, M.R., and Croteau, R. (1996) *J. Biol. Chem.* **271**:9201-9204).
- Monoterpene and diterpene biosynthesis are compartmentalized in plastids whereas sesquiterpene biosynthesis is cytosolic (reviewed in Kleinig, H. (1989) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **40**:39-53; Gershenzon, J., and Croteau, R. (1993) in *Lipid Metabolism in Plants* (Moore, T.S. Jr., ed) pp. 339-388, CRC Press, Boca Raton, FL; McGarvey, D.J., and Croteau, R. (1995) *Plant Cell* **7**, 1015-1026); thus, monoterpene and diterpene synthases are encoded as preproteins bearing an amino-terminal transit peptide for import of these nuclear gene products into plastids where they are proteolytically processed to the mature forms (Keegstra, K., Olsen, J.J., and Theg, S.M. (1989) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **40**:471-501; von Heijne, G., Stepphuhn, J., and Herrmann (1989) *Eur. J. Biochem.* **180**:535-545).
- Both the size of the deduced protein and the presence of an N-terminal domain (of 60 to 70 amino acids) with features characteristic of a targeting sequence [rich in serine residues (16-18%) and low in acidic residues (four Asp or Glu) (Keegstra, K., Olsen, J.J., and Theg, S.M. (1989) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **40**:471-501; von Heijne, G., Stepphuhn, J., and Herrmann (1989) *Eur. J. Biochem.* **180**:535-545)] suggest that AG2.2 (SEQ ID NO:1) encodes a monoterpene synthase rather than a sesquiterpene synthase or a diterpene synthase.

Since pAG2.2 contained the terpene synthase insert in reversed orientation, the ORF was subcloned in frame with glutathione S-transferase, for ultimate ease of purification (Bohlmann, J., DeLuca, V., Eilert, U., and Martin, W. (1995) *Plant J.* **7**:491-501; Bohlmann, J., Lins, T., Martin, W., and Eilert, U. (1996) *Plant Physiol.*

111:507-514), into pGEX-4T-2, yielding plasmid pGAG2.2. The recombinant fusion protein was expressed in *E. coli* strain XL1-Blue/pGAG2.2, then extracted and assayed for monoterpene synthase, sesquiterpene synthase and diterpene synthase activity using tritium labeled geranyl diphosphate, farnesyl diphosphate and geranylgeranyl diphosphate as the respective substrate. Enzymatic production of a terpene olefin was observed only with geranyl diphosphate as substrate, and the only product was shown to be myrcene by radio-GLC and GLC-MS comparison to an authentic standard (FIGURE 3). Bacteria transformed with pGEX vector containing the AG2.2 insert (SEQ ID NO:1) in antisense orientation did not afford detectable myrcene synthase activity when induced, and the protein isolated and assayed as above. A myrcene synthase cDNA has not been obtained previously from any source, although myrcene is a minor co-product (2%) of the native and recombinant limonene synthase from spearmint (Rajaonarivony, J.I.M., Gershenzon, J., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **296**:49-57; Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* **268**:23016-23024) and of several enzymes from sage (Croteau, R., and Satterwhite, D.M. (1989) *J. Biol. Chem.* **264**:15309-15315). cDNA cloning and functional expression of myrcene synthase, which is one of several wound-inducible monoterpene synthase activities of Grand fir (Gijzen, M., Lewinsohn, E., and Croteau, R. (1991) *Arch. Biochem. Biophys.* **289**:267-273), demonstrates that this acyclic monoterpene is formed by a distinct enzyme and is not a co-product of another synthase.

EXAMPLE 7

Cloning and Characterization of Clones AG3.18 (SEQ ID NO:3) Encoding (-)-Pinene Synthase and cDNA Clone AG10 (SEQ ID NO:5) Encoding

(-)-Limonene Synthase

Identification of cDNA Clone AG3.18 (SEQ ID NO:3) as (-)-Pinene Synthase and cDNA Clone AG10 (SEQ ID NO:5) as (-)-Limonene Synthase – Alignment of the four new terpene synthase cDNA sequences (AG1.28 (SEQ ID NO:15), AG2.2 (SEQ ID NO:1), AG4.30 (SEQ ID NO:17) and AG5.9 (SEQ ID NO:19)), and that for abietadiene synthase (Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268), allowed the identification of several conserved sequence motifs among this enzyme family from Grand fir, which provided the foundation for an extended similarity-based cloning approach. Two new sense primers E (SEQ ID NO:21) and F (SEQ ID NO:22) were designed according to conserved sequence elements, whereas a degenerate antisense

primer G (SEQ ID NO:23) was designed based upon very limited amino acid sequence information from pinene synthase (see Example 1). Only the combination of primers E (SEQ ID NO:21) and G (SEQ ID NO:23) amplified a specific product of 533 bps, which was designated as probe 3 (SEQ ID NO:24).

5 Hybridization of 10^5 Grand fir λ ZAP II cDNA clones with probe 3 (SEQ ID NO:24) yielded two types of signals comprised of about 400 strongly positive clones and an equal number of weak positives, indicating that the probe recognized more than one type of cDNA. Thirty-four of the former clones and 18 of the latter were purified, the inserts were selected by size (2.0-2.5 kb), and the *in vivo* excised clones
10 were partially sequenced from both ends. Those clones which afforded weak hybridization signals were shown to contain inserts that were either identical to myrcene synthase clone AG2.2 (SEQ ID NO:1) or exhibited no significant sequence similarity to terpene synthases. Clone AG3.48 (SEQ ID NO:31) contained the myrcene synthase ORF in the correct orientation and in frame for expression from
15 the Bluescript plasmid vector. This cDNA was functionally expressed in *E. coli* and the resulting enzyme was shown to accept only geranyl diphosphate as the prenyl diphosphate substrate and to produce myrcene as the exclusive reaction product. This finding with AG3.48 (SEQ ID NO:31) confirms that expression of AG2.2 (SEQ ID NO:1) as the glutathione *S*-transferase fusion protein from pGAG2.2 does not
20 influence substrate utilization or product outcome of the myrcene synthase.

Clones which gave strong hybridization signals segregated into distinct sequence groups represented by clone AG3.18 (SEQ ID NO:3)(2018 bp insert with ORF of 1884 nt; encoded protein of 628 residues at 71,505 Da and pI of 5.5) and AG10 (SEQ ID NO:5)(2089 bp insert with ORF of 1911 nt; encoded protein of 637
25 residues at 73,477 Da and pI of 6.4). AG3.18 (SEQ ID NO:3) and AG10 (SEQ ID NO:5) form a subfamily together with the myrcene synthase clone AG2.2 (SEQ ID NO:1) that is characterized by a minimum of 79% pairwise similarity (64% identity) at the amino acid level. Like myrcene synthase, both AG3.18 (SEQ ID NO:3) and AG10 (SEQ ID NO:5) encode *N*-terminal sequences of 60 to 70 amino acids which
30 are rich in serine (19-22% and 11-15%, respectively) and low in acidic residues (4 and 2, respectively) characteristic of plastid transit peptides (Keegstra, K., Olsen, J.J., and Theg, S.M. (1989) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **40**:471-501; von Heijne, G., Stepphuhn, J., and Herrmann (1989) *Eur. J. Biochem.* **180**:535-545).

Plasmid pAG3.18 (SEQ ID NO:3) contained the presumptive terpene
35 synthase ORF in frame for direct expression from the bluescript plasmid, whereas the

AG10 (SEQ ID NO:5) ORF was in reversed orientation. Both AG3.18 (SEQ ID NO:3) and AG10 (SEQ ID NO:5) were subcloned into expression vectors yielding plasmids pGAG3.18 and pSBAG10. Recombinant proteins were expressed in bacterial strain *E. coli* XL0LR/pAG3.18, *E. coli* XL1-Blue/pGAG3.18 and *E. coli* BL21(DE3)/pSBAG10. When extracts of the induced cells were tested for terpene synthase activity with all of the potential prenyl diphosphate substrates, only geranyl diphosphate was utilized. Extracts from *E. coli* BL21(DE3)/pSBAG10 converted geranyl diphosphate to limonene as the major product with lesser amounts of α -pinene, β -pinene and β -phellandrene, as determined by radio-GLC and combined GLC-MS (FIGURE 5). Chiral phase capillary GLC on β -cyclodextrin revealed the limonene product to be the (-)-4*S*-enantiomer and the pinene products to be the related (-)-(1*S*:5*S*)-enantiomers. Although optically pure standards were not available for the analysis, stereochemical considerations suggest that the minor product β -phellandrene is also the mechanistically related (-)-(4*S*)-antipode (Gambliel, H., and Croteau, R. (1984) *J. Biol. Chem.* **259**:740-748; Croteau, R., Satterwhite, D.M., Cane, D.E., and Chang, C.C. (1988) *J. Biol. Chem.* **263**:10063-10071; Wagschal, K., Savage, T.J., and Croteau, R. (1991) *Tetrahedron* **47**:5933-5944; Croteau, R., Satterwhite, D.M., Wheeler, C.J., and Felton, N.M. (1989) *J. Biol. Chem.* **264**:2075-2080; LaFever, R.E., and Croteau, R. (1993) *Arch. Biochem. Biophys.* **301**:361-366). Similar analysis of the monoterpene products generated from geranyl diphosphate by cell-free extracts of *E. coli* XL0LR/pAG3.18 and *E. coli* XL1-Blue/pGAG3.18 demonstrated the presence of a 42:58% mixture of α -pinene and β -pinene (FIGURE 4), the same product ratio previously described for the purified, native (-)-pinene synthase from Grand fir (Lewinsohn, E., Gijzen, M., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **293**:167-173). Chiral phase capillary GLC confirmed the products of the recombinant pinene synthase to be the (-)-(1*S*:5*S*)-enantiomers, as expected. No other monoterpene co-products were detected with the recombinant (-)-pinene synthase, as observed previously for the native enzyme (Lewinsohn, E., Gijzen, M., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **293**:167-173).

Evidence for the formation of both α - and β -pinene by a single enzyme has been previously provided through co-purification studies, and differential inhibition and inactivation studies, as well as by isotopically sensitive branching experiments (Gambliel, H., and Croteau, R. (1984) *J. Biol. Chem.* **259**:740-748; Wagschal, K.C., Pyun, H.-J., Coates, R.M., and Croteau, R. (1994) *Arch. Biochem. Biophys.* **308**:477-

487; Wagschal, K., Savage, T.J., and Croteau, R. (1991) *Tetrahedron* **47**:5933-5944; Croteau, R., Wheeler, C.J., Cane, D.E., Ebert, R., and Ha, H.-J. (1987) *Biochemistry* **26**:5383-5389). The cDNA cloning of pinene synthase provides the ultimate proof that a single enzyme forms both products. The calculated molecular weight of the (-)-pinene synthase deduced from *AG3.18* (SEQ ID NO:3) is approximately 64,000 (excluding the putative transit peptide), which agrees well with the molecular weight of 63,000 established for the native enzyme from Grand fir by gel permeation chromatography and SDS-PAGE (Lewinsohn, E., Gijzen, M., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **293**:167-173).

10 A limonene synthase cDNA has thus far been cloned only from two very closely related angiosperm species (Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* **268**:23016-23024; Yuba, A., Yazaki, K., Tabata, M., Honda, G., and Croteau, R. (1996) *Arch. Biochem. Biophys.* **332**:280-287), and the isolation of a pinene synthase cDNA has not been reported before. Pinene synthase has previously received considerable attention as a major defense-related monoterpene synthase in conifers (Gijzen, M., Lewinsohn, E., and Croteau, R. (1991) *Arch. Biochem. Biophys.* **289**:267-273; Lewinsohn, E., Gijzen, M., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **293**:167-173). In the Grand fir cDNA library, which was synthesized from mRNA obtained from wound-induced sapling stems, clones corresponding to pinene synthase are at least ten times more abundant than clones for myrcene synthase. This finding reflects the relative proportions of the induced levels of activities of these enzymes in Grand fir saplings; pinene synthase and limonene synthase are the major monoterpene synthase activities whereas the induced level of myrcene synthase activity is relatively low (Gijzen, M., Lewinsohn, E., and Croteau, R. (1991) *Arch. Biochem. Biophys.* **289**:267-273). The cDNAs for inducible monoterpene synthases provide probes for genetic and molecular analysis of oleoresin-based defense in conifers. Northern blots (FIGURE 6) of total RNA extracted from non-wounded sapling stems and from stems two days after wounding (when enzyme activity first appears) were probed with cDNA fragments for *AG2.2* (SEQ ID NO:1), *AG3.18* (SEQ ID NO:3) and *AG10* (SEQ ID NO:5), and thus demonstrated that increased mRNA accumulation for monoterpene synthases is responsible for this induced, defensive response in Grand fir. The availability of cloned, defense-related monoterpene synthases presents several possible avenues for transgenic manipulation of oleoresin composition to improve tree resistance to bark beetles and other pests. For example, altering the

- monoterpene content of oleoresin may chemically disguise the host and decrease insect aggregation by changing the levels of pheromone precursors or predator attractants, or lower infestation by increasing toxicity toward beetles and their pathogenic fungal associates (Johnson, M.A., and Croteau, R. (1987) in *Ecology and Metabolism of Plant Lipids* (Fuller, G., and Nes, W.D., eds) pp. 76-91, American Chemical Society Symposium Series 325, Washington, DC; Gijzen, M., Lewinsohn, E., Savage, T.J., and Croteau, R.B. (1993) in *Bioactive Volatile Compounds from Plants* (Teranishi, R., Buttery, R.G., and Sugisawa, H., eds) pp. 8-22, American Chemical Society Symposium Series 525, Washington, DC; Byers, J.A. (1995) in *Chemical Ecology of Insects 2* (Cardé, R.T., and Bell, W.J., eds) pp. 154-213, Chapman and Hall, New York).

EXAMPLE 8

Properties of the Recombinant Monoterpene Synthases Encoded by cDNA Clones AG2.2 (SEQ ID NO:1), AG3.18 (SEQ ID NO:3) and AG10 (SEQ ID NO:5)

- All three recombinant enzymes require Mn^{2+} for activity, and Mg^{2+} is essentially ineffective as the divalent metal ion cofactor. This finding confirms earlier results obtained with the native monoterpene synthases of Grand fir and lodgepole pine (*Pinus contorta*) (Lewinsohn, E., Gijzen, M., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **293**:167-173; Savage, T.J., Hatch, M.W., and Croteau, R. (1994) *J. Biol. Chem.* **269**:4012-4020). All terpene synthases and prenyltransferases are thought to employ a divalent metal ion, usually Mg^{2+} or Mn^{2+} , in the ionization steps of the reaction sequence to neutralize the negative charge of the diphosphate leaving group (Croteau, R. (1987) *Chem. Rev.* **87**:929-954; Cane, D.E. (1992) *Ciba Found. Symp. Ser.* **171**:163-167; Poulter, C.D., and Rilling, H.C. (1981) in *Biosynthesis of Isoprenoid Compounds* (Porter, J.W., and Spurgeon, S.L., eds) Vol. 1, pp. 161-224, Wiley & Sons, New York), and all relevant sequences thus far obtained bear a conserved aspartate rich element (DDXXD)(SEQ ID NO:45) considered to be involved in divalent metal ion binding (Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268; Ashby, M.N., and Edwards, P.A. (1990) *J. Biol. Chem.* **265**:13157-13164; Chen, A., Kroon, P.A., and Poulter, D.C. (1994) *Protein Sci.* **3**:600-607; Tarshis, L.C., Yan, M., Poulter, C.D., and Sacchettini, J.C. (1994) *Biochemistry* **33**:10871-10877; Cane, D.E., Sohng, J.K., Lamberson, C.R., Rudnicki, S.M., Wu, Z., Lloyd, M.D., Oliver, J.S., and Hubbard, B.R. (1994) *Biochemistry* **33**:5846-5857; Proctor, R.H., and

Hohn, T.M. (1993) *J. Biol. Chem.* **268**:4543-4548). In addition to this strict, general dependence on a divalent metal ion, the monoterpene synthases of conifers are unique in their further requirement for a monovalent cation (K^+), a feature that distinguishes the gymnosperm monoterpene synthases from their counterparts from angiosperm species and implies a fundamental structural and/or mechanistic difference between these two families of catalysts (Savage, T.J., Hatch, M.W., and Croteau, R. (1994) *J. Biol. Chem.* **269**:4012-4020). All three recombinant monoterpene synthases depend upon K^+ , with maximum activity achieved at approximately 500 mM KCl. A requirement for K^+ has been reported for a number of different types of enzymes, including those that catalyze phosphoryl cleavage or transfer reactions (Suelter, C.H. (1970) *Science* **168**:789-794) such as Hsc70 ATPase (Wilbanks, S.M., and McKay, D.B. (1995) *J. Biol. Chem.* **270**:2251-2257). The crystal structure of bovine Hsc70 ATPase indicates that both Mg^{2+} and K^+ interact directly with phosphate groups of the substrate and implicates three active site aspartate residues in Mg^{2+} and K^+ binding (Wilbanks, S.M., and McKay, D.B. (1995) *J. Biol. Chem.* **270**:2251-2257), reminiscent of the proposed role of the conserved DDXXD (SEQ ID NO:45) motif of the terpene synthases and prenyltransferases in divalent cation binding, a function also supported by recent site directed mutagenesis (Marrero, P.F., Poulter, C.D., and Edwards, P.A. (1992) *J. Biol. Chem.* **267**:21873-21878; Joly, A., and Edwards, P.A. (1993) *J. Biol. Chem.* **268**: 26983-26989; Song, L., and Poulter, C.D. (1994) *Proc. Natl. Acad. Sci. U.S.A.* **91**:3044-3048; Koyama, T., Tajima, M., Sano, H., Doi, T., Koike-Takeshita, A., Obata, S., Nishino, T., and Ogura, K. (1996) *Biochemistry* **35**:9533-9538) and by X-ray structural analysis (Tarshis, L.C., Yan, M., Poulter, C.D., and Sacchettini, J.C. (1994) *Biochemistry* **33**:10871-10877) of farnesyl diphosphate synthase.

cDNA cloning and functional expression of the myrcene, limonene and pinene synthases from Grand fir represent the first example of the isolation of multiple synthase genes from the same species, and provide tools for evaluation of structure-function relationships in the construction of acyclic, monocyclic and bicyclic monoterpene products and for detailed comparison to catalysts from phylogenetically distant plants that carry out ostensibly identical reactions (Gambliel, H., and Croteau, R. (1984) *J. Biol. Chem.* **259**:740-748; Rajaonarivony, J.I.M., Gershenzon, J., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **296**:49-57; Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* **268**:23016-23024; Adam, K.-P., Crock, J., and Croteau, R. (1996) *Arch.*

- Biochem. Biophys.* **332**:352-356). The recent acquisition of cDNA isolates encoding sesquiterpene synthases and diterpene synthases (Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268) from Grand fir should, together with the monoterpene synthases, also permit addressing the structural basis of chain-length specificity for prenyl diphosphate substrates in this family of related enzymes.

EXAMPLE 9

Sequence Comparison of Cloned Monoterpene Synthases

- Previous studies based on substrate protection from inactivation with selective amino acid modifying reagents have implicated functionally important cysteine, histidine and arginine residues in a range of different monoterpene synthases (Rajaonarivony, J.I.M., Gershenzon, J., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **296**:49-57; Lewinsohn, E., Gijzen, M., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **293**:167-173; Savage, T.J., Hatch, M.W., and Croteau, R. (1994) *J. Biol. Chem.* **269**:4012-4020; Rajaonarivony, J.I.M., Gershenzon, J., Miyazaki, J., and Croteau, R. (1992) *Arch. Biochem. Biophys.* **299**:77-82; Savage, T.J., Ichii, H., Hume, S.D., Little, D.B., and Croteau, R. (1994) *Arch. Biochem. Biophys.* **320**:257-265). Sequence alignment of 21 terpene synthases of plant origin (Colby, S.M., Alonso, W.R., Katahira, E.J., McGarvey, D.J., and Croteau, R. (1993) *J. Biol. Chem.* **268**:23016-23024; Stofer Vogel, B., Wildung, M.R., Vogel, G., and Croteau, R. (1996) *J. Biol. Chem.* **271**:23262-23268; Facchini, P.J., and Chappell, J. (1992) *Proc. Natl. Acad. Sci. USA* **89**:11088-11092; Mau, C.J.D., and West, C.A. (1994) *Proc. Natl. Acad. Sci. USA* **91**:8497-8501; Yuba, A., Yazaki, K., Tabata, M., Honda, G., and Croteau, R. (1996) *Arch. Biochem. Biophys.* **332**:280-287; Wildung, M.R., and Croteau, R. (1996) *J. Biol. Chem.* **271**:9201-9204; Yamaguchi, S., Saito, T., Abe, H., Yamane, H., Murofushi, N., and Kamiya, Y. (1996) *Plant J.* **10**:203-213; Dudareva, N., Cseke, L., Blanc, V.M., and Pichersky, E. (1996) *Plant Cell* **8**:1137-1148; Chen, X.-Y., Chen, Y., Heinsteins, P., and Davisson, V.J. (1995) *Arch. Biochem. Biophys.* **324**:255-266; Chen, X.-Y., Wang, M., Chen, Y., Davisson, J., and Heinsteins, P. (1996) *J. Nat. Prod.* **59**:944-951; Back, K., and Chappell, J. (1995) *J. Biol. Chem.* **270**:7375-7381) reveals two absolutely conserved arginine residues, corresponding to Arg¹⁸⁴ and Arg³⁶⁵ of pinene synthase (SEQ ID NO:4), one highly conserved cysteine residue (pinene synthase Cys⁵⁴³)(SEQ ID NO:4), and one highly conserved histidine residue (pinene synthase His¹⁸⁶)(SEQ ID NO:4). The DDXXD (SEQ ID NO:45) sequence motif (pinene synthase Asp³⁷⁹, Asp³⁸⁰ and Asp³⁸³) (SEQ

ID NO:4) is absolutely conserved in all relevant plant terpene synthases, as are several other amino acid residues corresponding to Phe¹⁹⁸, Leu²⁴⁸, Glu³²², Trp³²⁹, Trp⁴⁶⁰ and Pro⁴⁶⁷ of pinene synthase (SEQ ID NO:4).

5 Amino acid sequences of the plant terpene synthases were compared with each other and with the deduced sequences of several sesquiterpene synthases cloned from microorganisms (Proctor, R.H., and Hohn, T.M. (1993) *J. Biol. Chem.* **268**:4543-4548; Back, K., and Chappell, J. (1995) *J. Biol. Chem.* **270**:7375-7381; Hohn, T.M., and Desjardins, A.E. (1992) *Mol. Plant-Microbe Interactions* **5**:249-256). As with all other plant terpene synthases, no significant
10 conservation in primary sequence exists between the monoterpene synthases from Grand fir and the terpene synthases of microbial origin, except for the DDXXD (SEQ ID NO:45) sequence motif previously identified as a common element of all terpene synthases, and prenyltransferases which employ a related electrophilic reaction mechanism (Croteau, R., Wheeler, C.J., Cane, D.E., Ebert, R., and Ha, H.-J. (1987) *Biochemistry* **26**:5383-5389; Chen, A., Kroon, P.A., and Poulter, D.C. (1994) *Protein Sci.* **3**:600-607; McCaskill, D., and Croteau, R. (1997) *Adv. Biochem. Engineering Biotech.* **55**:108-146). The evidence is presently insufficient to determine whether
15 extant plant and microbial terpene synthases represent divergent evolution from a common ancestor, which may also have given rise to the prenyltransferases, or whether these similar catalysts evolved convergently.
20

EXAMPLE 10

Alteration of Monoterpene Levels and Composition in Plant Seeds

In accordance with the present invention, methods for increasing production of monoterpene compounds in a plant, particularly in plant seeds, are provided. The
25 methods involve transforming a plant cell with a nucleic acid sequence encoding at least one gymnosperm monoterpene synthase, such as those encoded by the nucleic acid sequences set forth in SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:5. This has the effect of altering monoterpene biosynthesis, thereby increasing the production of monoterpenes, as well as providing novel seed oils having desirable monoterpene
30 compositions. In this manner, the transformed seed provides a factory for the production of modified oils. The modified oil itself may be used and/or the compounds in the oils can be isolated. Thus, the present invention allows for the production of particular monoterpenes of interest as well as speciality oils.

The nucleic acid encoding the monoterpene synthases of the present invention
35 can be used in expression cassettes for expression in the transformed plant tissues.

To alter the monoterpene levels in a plant of interest, the plant is transformed with at least one expression cassette comprising a transcriptional initiation region linked to a nucleic acid sequence encoding a monoterpene synthase. Such an expression cassette is provided with a plurality of restriction sites for insertion of the nucleic acid sequence encoding a monoterpene synthase so that it is under the transcriptional regulation of the regulatory regions.

The transcriptional initiation sequence may be native or analogous to the host or foreign or heterologous to the host. In this regard, the term "foreign" means that the transcriptional initiation sequence is not found in the wild-type host into which the transcriptional initiation region is introduced.

Of particular interest are those transcriptional initiation regions associated with storage proteins, such as napin, cruciferin, β -conglycinin, phaseolin, globulin or the like, and proteins involved in fatty acid biosynthesis, such as acyl carrier protein (ACP). See, U.S. Patent No. 5,420,034, herein incorporated by reference.

The transcriptional cassette will preferably include, in the 5' to 3' direction of transcription, a transcriptional and translational initiation region, a gymnosperm monoterpene synthase DNA sequence of interest, and a transcriptional and translational termination region functional in plants. The termination region may be from the same organism as the transcriptional initiation region, may be from the same organism as the monoterpene synthase DNA, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. Other termination sequences are set forth in Guerineau et al., (1991), *Mol. Gen. Genet.*, **262**:141-144; Proudfoot, (1991), *Cell*, **64**:671-674; Sanfacon et al., (1991), *Genes Dev.*, **5**:141-149; Mogen et al., (1990), *Plant Cell*, **2**:1261-1272; Munroe et al., (1990), *Gene*, **91**:151-158; Ballas et al., (1989), *Nucleic Acids Res.*, **17**:7891-7903; Joshi et al., (1987), *Nucleic Acid Res.*, **15**:9627-9639).

In the presently preferred form of the invention, a nucleic acid sequence encoding a gymnosperm monoterpene synthase protein will be targeted to plastids, such as chloroplasts, for expression. Thus, the nucleic acid sequence, or sequences, encoding a gymnosperm monoterpene synthase protein, or proteins, may be inserted into the plastid for expression with appropriate plastid constructs and regulatory elements. Alternatively, nuclear transformation may be used in which case the expression cassette will contain a nucleic acid sequence encoding a transit peptide to direct the monoterpene biosynthesis enzyme of interest to the plastid. Such transit

peptides are known in the art. See, for example, Von Heijne et al. (1991) *Plant Mol. Biol. Rep.* **9**:104-126; Clark et al. (1989) *J. Biol. Chem.* **264**:17544-17550; della-Cioppa et al. (1987) *Plant Physiol.* **84**:965-968; Romer et al. (1993) *Biochem. Biophys. Res. Commun.* **196**:1414-1421; and, Shah et al. (1986) *Science* **233**:478-481. Nucleic acid sequences encoding gymnosperm monoterpene synthases of the present invention may utilize native or heterologous transit peptides.

The construct may also include any other necessary regulators such as plant translational consensus sequences (Joshi, C.P., (1987), *Nucleic Acids Research*, **15**:6643-6653), introns (Luehrsen and Walbot, (1991), *Mol. Gen. Genet.*, **225**:81-93) and the like, operably linked to a nucleotide sequence encoding a monoterpene synthase of the present invention.

It may be beneficial to include 5' leader sequences in the expression cassette which can act to enhance translation. Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5' noncoding region) (Elroy-Stein, O., Fuerst, T.R., and Moss, B. (1989) *PNAS USA* **86**:6126-6130); potyvirus leaders, for example, TEV leader (Tobacco Etch Virus) (Allison et al. (1986); MDMV leader (Maize Dwarf Mosaic Virus); *Virology*, **154**:9-20), and human immunoglobulin heavy-chain binding protein (BiP), (Macejak, D.G., and Sarnow, P. (1991), *Nature*, **353**:90-94; untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4), (Jobling, S.A., and Gehrke, L. (1987), *Nature*, **325**:622-625; tobacco mosaic virus leader (TMV), (Gallie, D.R. et al. (1989), *Molecular Biology of RNA*, pages 237-256; and maize chlorotic mottle virus leader (MCMV) (Lommel, S.A. et al. (1991), *Virology*, **81**:382-385. See also, Della-Cioppa et al., (1987), *Plant Physiology*, **84**:965-968.

Depending upon where the monoterpene synthase sequence of interest is to be expressed, it may be desirable to synthesize the sequence with plant preferred codons, or alternatively with chloroplast preferred codons. The plant preferred codons may be determined from the codons of highest frequency in the proteins expressed in the largest amount in the particular plant species of interest. See, EPA 0359472; EPA 0385962; WO 91/16432; Perlak et al. (1991) *Proc. Natl. Acad. Sci. USA* **88**:3324-3328; and Murray et al. (1989) *Nucleic Acids Research* **17**:477-498. In this manner, the nucleotide sequences can be optimized for expression in any plant. It is recognized that all or any part of the nucleic acid sequence encoding a gymnosperm monoterpene synthase protein may be optimized or synthetic. That is,

synthetic or partially optimized sequences may also be used. For the construction of chloroplast preferred genes, see U.S. Patent No. 5,545,817.

5 In preparing the transcription cassette, the various DNA fragments may be manipulated, so as to provide for the DNA sequences in the proper orientation and in the proper reading frame. Towards this end, adapters or linkers may be employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites, or the like. For this purpose, *in vitro* mutagenesis, primer repair, restriction, annealing, resection, ligation, or the like may be employed, where insertions, deletions or substitutions, such as transitions and transversions, may be involved.

10 The recombinant DNA molecules of the invention can be introduced into the plant cell in a number of art-recognized ways. Those skilled in the art will appreciate that the choice of method might depend on the type of plant, *i.e.*, monocot or dicot, targeted for transformation. Suitable methods of transforming plant cells include microinjection (Crossway et al. (1986) *BioTechniques* 4:320-334), electroporation (Riggs et al. (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606), *Agrobacterium* mediated transformation (Hinchee et al. (1988) *Biotechnology* 6:915-921) and ballistic particle acceleration (see, for example, Sanford et al., U.S. Patent No. 4,945,050; and McCabe et al. (1988) *Biotechnology* 6:923-926). Also see, Weissinger et al. (1988) *Annual Rev. Genet.* 22:421-477; Sanford et al. (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou et al. (1988) *Plant Physiol.* 87:671-674 (soybean); McCabe et al. (1988) *Bio/Technology* 6:923-926 (soybean); Datta et al. (1990) *Biotechnology* 8:736-740 (rice); Klein et al. (1988) *Proc. Natl. Acad. Sci. USA*, 85:4305-4309 (maize); Klein et al. (1988) *Biotechnology* 6:559-563 (maize); Klein et al. (1988) *Plant Physiol.* 91:440-444 (maize); Fronun et al. (1990) *Biotechnology* 8:833-839; and Gordon-Kamm et al. (1990) *Plant Cell* 2:603-618 (maize).

20 Alternatively, a plant plastid can be transformed directly. Stable transformation of chloroplasts has been reported in higher plants, see, for example, SVAB et al. (1990) *Proc. Nat'l. Acad. Sci. USA* 87:85268530; SVAB & Maliga (1993) *Proc. Natl. Acad. Sci. USA* 90:913-917; Staub & Maliga (1993) *Embo J.* 12:601-606. The method relies on particle gun delivery of DNA containing a selectable marker and targeting of the DNA to the plastid genome through homologous recombination. In such methods, plastid gene expression can be accomplished by use of a plastid gene promoter or by trans-activation of a silent

35

plastid-borne transgene positioned for expression from a selective promoter sequence such as that recognized by T7 RNA polymerase. The silent plastid gene is activated by expression of the specific RNA polymerase from a nuclear expression construct and targeting of the polymerase to the plastid by use of a transit peptide. Tissue-specific expression may be obtained in such a method by use of a nuclear-encoded and plastid-directed specific RNA polymerase expressed from a suitable plant tissue specific promoter. Such a system has been reported in McBride et al. (1994) *Proc. Natl. Acad. Sci. USA* **91**:7301-7305.

The cells which have been transformed may be grown into plants by a variety of art-recognized means. See, for example, McConnick et al., *Plant Cell Reports* (1986), **5**:81-84. These plants may then be grown, and either selfed or crossed with a different plant strain, and the resulting homozygotes or hybrids having the desired phenotypic characteristic identified. Two or more generations may be grown to ensure that the subject phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure the desired phenotype or other property has been achieved.

As a host cell, any plant variety may be employed. Of particular interest, are plant species which provide seeds of commercial value. For the most part, plants will be chosen where the seed is produced in high amounts, a seed-specific product of interest is involved, or the seed or a seed part is edible. Seeds of interest in the practice of the present invention include, but are not limited to, the oil seeds, such as oilseed Brassica seeds, cotton seeds, soybean, safflower, sunflower, coconut, palm, and the like; grain seeds such as wheat, barley, oats, amaranth, flax, rye, triticale, rice and corn; other edible seeds or seeds with edible parts including pumpkin, squash, sesame, poppy, grape, mung beans, peanut, peas, beans, radish, alfalfa, cocoa, and coffee; and tree nuts such as walnuts, almonds, pecans, and chick-peas.

EXAMPLE 11

A Strategy For Cloning Gymnosperm Monoterpene Synthases

The present invention includes gymnosperm monoterpene synthase proteins, and nucleic acid molecules that encode gymnosperm monoterpene synthase proteins. The amino acid sequence of each of the gymnosperm monoterpene synthase proteins of the present invention each includes at least one of the amino acid sequence elements disclosed in Table 1.

Table 1

Amino Acids	Sequence	Orientation	Comments
-------------	----------	-------------	----------

1. 70-77	H S N (L, I, V) W D D D (SEQ ID NO: 46)	F only	HS makes a poor reverse primer
2. 148-153	A L D Y V Y (SEQ ID NO: 47)	F and R	
3. 306-312	E L A K L E F (SEQ ID NO: 48)	F and R	
4. 328-333	R W W K E S (SEQ ID NO: 49)	F and R	F primer uses 1 st nt of Ser codon; R uses 1 st two nts of Arg codon (rare only)
5. 377-383	(V, I, L) L D D M Y D (SEQ ID NO: 50)	F and R	
6. 377-383	(V, I, L) L D D L Y D (SEQ ID NO: 51)	F and R	Degeneracy of M/L at 381 too high for single primer
7. 377-383	(V, I, L) L D D I Y D (SEQ ID NO: 52)	F and R	
8. 543-549	C Y M K D (N, H) P (SEQ ID NO: 53)	R	F primer can also be constructed with this peptide but is too close to the 3' end to be useful

The numbers set forth in Table 1 for the first and last amino acid residue of each of the peptide sequences is the number of the corresponding amino acid residue in the amino acid sequence of the (-)-pinene synthase (SEQ ID NO:4) isolated from *Abies grandis*. Where a sequence of amino acid residues appears in brackets, e.g., (L,I,V) in Table 1, the first amino acid residue within the brackets is the residue that appears in the (-)-pinene synthase amino acid sequence set forth in SEQ ID NO:4. The subsequent amino acid residues within the brackets represent other amino acid residues that commonly occur at the corresponding position in the amino acid sequence of other *Abies grandis* enzymes involved in terpene synthesis.

In Table 1, the letter "F" refers to the forward PCR reaction, i.e., the PCR reaction which synthesizes the sense nucleic acid strand that encodes a gymnosperm monoterpene synthase. The letter "R" refers to the reverse PCR reaction, i.e., the PCR reaction that synthesizes the antisense nucleic acid molecule that is complementary to the sense nucleic acid strand synthesized in the forward PCR reaction.

In order to clone nucleic acid molecules encoding gymnosperm monoterpene synthases of the present invention, one or more oligonucleotide molecules corresponding to at least a portion of one of the amino acid sequences set forth in Table 1 can be used as a probe or probes with which to screen a genomic or cDNA library derived from one or more gymnosperm species. In this context, the term "corresponding," or "correspond" or "corresponds," means that the oligonucleotide base sequence either a) encodes all or part of at least one of the amino acid sequences set forth in Table 1, or b) is complementary to a base sequence that encodes all or part of at least one of the amino acid sequences set forth in Table 1. The oligonucleotide probe(s) may contain a synthetic base, such as inosine, which can be substituted for one or more of the four, naturally-occurring bases, *i.e.*, adenine ("A"), guanine ("G"), cytosine ("C") and thymine ("T"). Thus, for example, the following oligonucleotide sequences "correspond" to the tripeptide sequence M M M: 5'ATGATGATG3' (sense orientation) (SEQ ID NO:54); 3'TACTACTAC5' (antisense orientation) (SEQ ID NO:55) and 3'IACIACIAC5' (SEQ ID NO:56).

One or more oligonucleotide sequence(s), corresponding to at least a portion of at least one of the amino acid sequences set forth in Table 1, can be used to screen a nucleic acid library in order to identify monoterpene synthase clones of the present invention, according to methods well known to one of ordinary skill in the art. *See, e.g., Sambrook et al, supra.* The stringency of the hybridization and wash conditions during library screening in accordance with the present invention, utilizing one or more oligonucleotide sequence(s) corresponding to at least a portion of at least one of the amino acid sequences set forth in Table 1, is at least: for the hybridization step, 6X SSPE, 40-45°C, for 36 hours; for the wash step, 3X SSPE, 45°C, 3 X 15 minute washes. The presently preferred hybridization and wash conditions during library screening, utilizing one or more oligonucleotide sequence(s) corresponding to at least a portion of at least one of the amino acid sequences set forth in Table 1, in accordance with the present invention are: for the hybridization step, 6X SSPE, 40-45°C, for 36 hours; for the wash step, 0.1X SSPE, 65°C-70°C, 3 X 15 minute washes.

Examples of oligonucleotide sequences, corresponding to at least one of the amino acid sequences set forth in Table 1, that hybridize, under the foregoing hybridization and wash conditions, to the sense strands of the nucleic acid sequences of the present invention that encode gymnosperm monoterpene synthase proteins are set forth in Table 2.

Table 2

Nucleic Acid Sequence	Corresponds to:
GTG TCG TTG GAG ACC CTG CTG CTG (SEQ ID NO:57)	SEQ ID No. 46
CGG GAG CTG ATG CAG ATG (SEQ ID NO:58)	SEQ ID No. 47
CTC GAG CGG TTC GAG CTC AAG (SEQ ID NO:59)	SEQ ID No. 48
GCC ACC ACC TTC CTC TCG (SEQ ID NO:60)	SEQ ID No. 49
GAG GAG CTG CTG TAC ATG CTG (SEQ ID NO:61)	SEQ ID No. 50
GAG GAG CTG CTG GAG ATG CTG (SEQ ID NO:62)	SEQ ID No. 51

Similarly, each of the gymnosperm monoterpene synthase clones set forth in SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:5, or a portion thereof, may be used as a probe to screen a nucleic acid library in order to isolate monoterpene synthase clones of the present invention, according to methods well known to one of ordinary skill in the art. *See, e.g., Sambrook et al, supra.* The stringency of the hybridization and wash conditions during library screening in accordance with the present invention is at least: for the hybridization step, 6X SSPE buffer at 45°C to 50°C for 36 hours; for the wash step, 3X SSPE buffer at 50°C (three, fifteen minute washes).

In accordance with the present invention, the presently preferred hybridization and wash conditions during library screening utilizing any of the gymnosperm monoterpene synthase clones set forth in SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:5, or a portion thereof, as probe are: for the hybridization step, 6X SSPE, 40-45°C, for 36 hours; for the wash step, 0.1X SSPE, 70°C-75°C, 3 X 15 minute washes.

Additionally, at least two oligonucleotide sequence(s), each corresponding to at least a portion of at least one of the amino acid sequences set forth in Table 1, can be used in a PCR reaction to generate a portion of a monoterpene synthase clone of the present invention, which can be used as a probe to isolate a full-length clone of a monoterpene synthase clone of the present invention. Thus, oligonucleotides that are useful as probes in the forward PCR reaction correspond to at least a portion of at least one of the amino acid sequences disclosed in Table 1 as having the "F" orientation. Conversely, oligonucleotides that are useful as probes in the reverse

PCR reaction correspond to at least a portion of at least one of the amino acid sequences disclosed in Table 1 as having the "R" orientation. PCR reactions can be carried out according to art-recognized PCR reaction conditions, such as the PCR reaction conditions set forth in Example 1 herein and as set forth in "PCR Strategies",

- 5 M.A. Innis, D.H. Gelfand and J.J. Sninsky, eds., 1995, Academic Press, San Diego, CA (Chapter 14); "PCR Protocols: A Guide to Methods and Applications". M.A. Innis, D.H. Gelfand, J.J. Sninsky and T.J. White, eds., Academic Press, NY (1990). The presently preferred PCR reaction conditions are:

dNTPs	200 μ M each
MgCl ₂	5-7 mM
F and R primers	100 nM - 1 μ M each
Taq polymerase	1-2 units/reaction
cDNA template	10-100 ng/reaction
Buffers, PCR grade water, and Chill-out wax or mineral oil	

10

The presently preferred thermocycler conditions are:

Denaturation	94 ⁰ x 2 min	1 cycle
Denaturation	94 ⁰ x 45 s	35 cycles
Annealing	42 ⁰ - 55 ⁰ x 45 s - 1 min	"
Polymerization	72 ⁰ x 1-2 min	"
Adenylation	72 ⁰ x 10 min	1 cycle

- 15 While the preferred embodiment of the invention has been illustrated and described, it will be appreciated that various changes can be made therein without departing from the spirit and scope of the invention.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. An isolated nucleotide sequence encoding a gymnosperm monoterpene synthase, said isolated nucleotide sequence being capable of hybridizing under stringent conditions to at least one oligonucleotide having a base sequence that corresponds to at least one peptide sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.
2. An isolated nucleotide sequence of Claim 1 encoding a gymnosperm myrcene synthase.
3. An isolated nucleotide sequence of Claim 1 encoding a gymnosperm limonene synthase.
4. An isolated nucleotide sequence of Claim 1 encoding a gymnosperm pinene synthase.
5. An isolated nucleotide sequence of Claim 1 encoding a Grand fir (*Abies grandis*) monoterpene synthase.
6. An isolated nucleotide sequence of Claim 1 encoding a Grand fir (*Abies grandis*) myrcene synthase.
7. An isolated nucleotide sequence of Claim 1 encoding a Grand fir (*Abies grandis*) limonene synthase.
8. An isolated nucleotide sequence of Claim 1 encoding a Grand fir (*Abies grandis*) pinene synthase.
9. An isolated nucleotide sequence of Claim 1 having the sequence of SEQ ID NO:1.
10. An isolated nucleotide sequence of Claim 1 having the sequence of SEQ ID NO:3.

11. An isolated nucleotide sequence of Claim 1 having the sequence of SEQ ID NO:5

12. An isolated nucleotide sequence of Claim 1 which encodes the amino acid sequence of SEQ ID NO:2.

13. An isolated nucleotide sequence of Claim 1 which encodes the amino acid sequence of SEQ ID NO:4

14. An isolated nucleotide sequence of Claim 1 which encodes the amino acid sequence of SEQ ID NO:6.

15. An isolated nucleotide sequence encoding a gymnosperm monoterpene synthase protein, said isolated nucleotide sequence having a complementary nucleotide sequence that is capable of hybridizing to the nucleotide sequence of any one of SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:5.

16. An isolated nucleic acid molecule encoding a gymnosperm monoterpene synthase, said nucleic acid molecule being capable of hybridizing to at least two oligonucleotides, each of said oligonucleotides having a base sequence that corresponds to at least one peptide sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

17. An isolated nucleic acid molecule encoding a Grand fir (*Abies grandis*) monoterpene synthase, said nucleic acid molecule being capable of hybridizing to at least two oligonucleotides, each of said oligonucleotides having a base sequence that corresponds to at least one peptide sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

18. An isolated gymnosperm monoterpene synthase protein, said gymnosperm monoterpene synthase protein including at least one amino acid sequence element selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

19. An isolated gymnosperm myrcene synthase protein, said gymnosperm myrcene synthase protein including at least one amino acid sequence element selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

20. An isolated gymnosperm pinene synthase protein, said gymnosperm pinene synthase protein including at least one amino acid sequence element selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

21. An isolated gymnosperm limonene synthase protein, said gymnosperm limonene synthase protein including at least one amino acid sequence element selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

22. An isolated Grand fir (*Abies grandis*) monoterpene synthase protein, said Grand fir (*Abies grandis*) monoterpene synthase protein including at least one amino acid sequence element selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

23. An isolated Grand fir (*Abies grandis*) monoterpene synthase protein of Claim 22, said monoterpene synthase protein having the sequence of SEQ ID NO:2.

24. An isolated Grand fir (*Abies grandis*) monoterpene synthase protein of Claim 22, said monoterpene synthase protein having the sequence of SEQ ID NO:4.

25. An isolated Grand fir (*Abies grandis*) monoterpene synthase protein of Claim 22, said monoterpene synthase protein having the sequence of SEQ ID NO:6.

26. A replicable expression vector comprising a nucleotide sequence encoding a gymnosperm monoterpene synthase, said nucleotide sequence encoding a gymnosperm monoterpene synthase being capable of hybridizing at high stringency to at least one oligonucleotide having a base sequence that corresponds to at least one peptide sequence selected from the group consisting of SEQ ID NO:46, SEQ ID

NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

27. A replicable expression vector of Claim 26 wherein said nucleotide sequence encoding a gymnosperm monoterpene synthase encodes a Grand fir (*Abies grandis*) monoterpene synthase.

28. A host cell comprising a vector of any one of claims 26 and 27.

29. A replicable expression vector comprising a nucleotide sequence encoding a gymnosperm monoterpene synthase, said nucleotide sequence having a complementary nucleotide sequence that is capable of hybridizing to the nucleotide sequence of any one of SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:5.

30. A host cell comprising a vector of Claim 29.

31. A method of enhancing the production of a gymnosperm monoterpene synthase in a suitable host cell comprising introducing into the host cell an expression vector of Claim 26 under conditions enabling expression of the gymnosperm monoterpene synthase in the host cell.

32. A method for enhancing monoterpene levels in a host cell, said method comprising transforming said host cell with a nucleic acid construct comprising at least one nucleic acid sequence defining a transcriptional initiation region from a gene expressed in the host cell and at least one nucleic acid sequence encoding a gymnosperm monoterpene synthase, said nucleic acid sequence encoding a gymnosperm monoterpene synthase being capable of hybridizing at high stringency to at least one oligonucleotide having a base sequence that corresponds to at least one peptide sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

33. The method of Claim 32 wherein said nucleic acid sequence encoding a gymnosperm monoterpene synthase encodes a Grand fir (*Abies grandis*) monoterpene synthase.

34. A method for altering monoterpene levels in a seed from a host plant, said method comprising transforming said host plant with a nucleic acid construct

comprising at least one nucleic acid sequence defining a transcriptional initiation region from a gene expressed in a plant seed and at least one nucleic acid sequence encoding a gymnosperm monoterpene synthase, said nucleic acid sequence encoding a gymnosperm monoterpene synthase being capable of hybridizing at high stringency to at least one oligonucleotide having a base sequence that corresponds to at least one peptide sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

35. The method of Claim 34 wherein the transcriptional initiation region is from a gene preferentially expressed in a plant seed.

36. The method of Claim 34 wherein said nucleic acid sequence encoding a gymnosperm monoterpene synthase encodes a Grand fir (*Abies grandis*) monoterpene synthase.

37. The method of Claim 34 wherein said gymnosperm monoterpene synthase is not native to said host plant.

38. A transgenic plant produced according to the method of Claim 34.

39. A method for increasing the monoterpene biosynthetic flux in seed from a host plant, said method comprising transforming said host plant with a construct comprising as operably linked components, a transcriptional initiation region from a gene preferentially expressed in a plant seed, a plastid transit peptide, a DNA sequence encoding a gymnosperm monoterpene synthase protein, and a transcriptional termination region, said DNA sequence encoding a gymnosperm monoterpene synthase protein being capable of hybridizing at high stringency to at least one oligonucleotide having a base sequence that corresponds to at least one peptide sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

40. The method of Claim 39, wherein said DNA sequence encoding a gymnosperm monoterpene synthase protein encodes an enzyme selected from the group consisting of limonene synthase, myrcene synthase and pinene synthase.

41. A method for producing a monoterpene compound in a seed, said method comprising the steps of obtaining a transformed plant which produces said seed, said plant having and expressing in its genome:

a nucleic acid sequence encoding a gymnosperm monoterpene synthase protein which is operably linked to a plastid transit peptide and a transcriptional initiation region from a gene preferentially expressed in a plant seed, wherein said nucleic acid sequence encoding a gymnosperm monoterpene synthase protein is capable of hybridizing at high stringency to at least one oligonucleotide having a base sequence that corresponds to at least one peptide sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53.

42. A seed transformed with a nucleic acid sequence encoding a gymnosperm monoterpene synthase protein, said nucleic acid sequence encoding a gymnosperm monoterpene synthase protein being capable of hybridizing at high stringency to at least one oligonucleotide having a base sequence that corresponds to at least one peptide sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52 and SEQ ID NO:53, said transformed seed having altered monoterpene levels.

43. The transformed seed of claim 42, wherein said seed produces increased levels of at least one monoterpene compound, said monoterpene compound selected from the group consisting of (-)-4S-limonene, (-)- α -pinene, myrcene, 4S- β -phellandrene.

44. The transformed seed of claim 43, wherein said seed produces increased levels of (-)-4S-limonene.

45. Seed produced by the method of any one of claims 34, 35, 36, 37, 39, 40 and 41.

46. Plants produced by the method of any one of claims 34, 35, 36, 37, 39, 40 and 41.

47. Oil extracted from seeds produced by the method of any one of claims 34, 35, 36, 37, 39, 40 and 41.

48. Meal extracted from seed produced by the method of any one of claims 34, 35, 36, 37, 39, 40 and 41.

49. The method of any one of claims 34, 35, 36, 37, 39, 40 and 41, wherein said seed is from a plant selected from the group consisting of oilseed Brassica, cotton, soybean, safflower, sunflower, coconut, palm, wheat, barley, rice, corn, oats, amaranth, pumpkin, squash, sesame, poppy, grape, mung beans, peanut, peas, beans, radish, alfalfa, cocoa, coffee, and tree nuts.

50. The method of Claim 49, wherein said seed is from an oilseed crop plant selected from the group consisting of oilseed brassica, cotton, soybean, safflower, sunflower, palm, coconut, and corn.

1/7




Fig. 1

2/7




Fig. 2

3/7




Fig. 3A




Fig. 3B




Fig. 3C




Fig. 4A







Fig. 4B



6/7




Fig. 5A




Fig. 5B

7/7




Fig. 5C

SEQUENCE LISTING

<110> Croteau, Rodney B
 Bohlmann, Carl J
 Steele, Christopher L

<120> Monoterpene Synthases from Grand Fir (*Abies grandis*)

<130> WSUR112537

<140>

<141>

<150> 60/052,249

<151> 1997-07-11

<160> 62

<170> PatentIn Ver. 2.0

<210> 1

<211> 2196

<212> DNA

<213> *Abies grandis*

<220>

<221> CDS

<222> (69)..(1952)

<223> Clone AG2.2 encoding myrcene synthase

<400> 1

```

tgccggcacg aggttatctt gagcttcctc catataggcc aacacatata atatcaaagg 60
gagcaaga atg gct ctg gtt tct atc tca ccg ttg gct tcg aaa tct tgc 110
      Met Ala Leu Val Ser Ile Ser Pro Leu Ala Ser Lys Ser Cys
        1             5             10

ctg cgc aag tcg ttg atc agt tca att cat gaa cat aag cct ccc tat 158
Leu Arg Lys Ser Leu Ile Ser Ser Ile His Glu His Lys Pro Pro Tyr
  15             20             25             30

aga aca atc cca aat ctt gga atg cgt agg cga ggg aaa tct gtc acg 206
Arg Thr Ile Pro Asn Leu Gly Met Arg Arg Arg Gly Lys Ser Val Thr
      35             40             45

cct tcc atg agc atc agt ttg gcc acc gct gca cct gat gat ggt gta 254
Pro Ser Met Ser Ile Ser Leu Ala Thr Ala Ala Pro Asp Asp Gly Val
      50             55             60

caa aga cgc ata ggt gac tac cat tcc aat atc tgg gac gat gat ttc 302
Gln Arg Arg Ile Gly Asp Tyr His Ser Asn Ile Trp Asp Asp Asp Phe
      65             70             75

ata cag tct cta tca acg cct tat ggg gaa ccc tct tac cag gaa cgt 350
Ile Gln Ser Leu Ser Thr Pro Tyr Gly Glu Pro Ser Tyr Gln Glu Arg
      80             85             90

gct gag aga tta att gtg gag gta aag aag ata ttc aat tca atg tac 398
Ala Glu Arg Leu Ile Val Glu Val Lys Lys Ile Phe Asn Ser Met Tyr
      95             100             105             110

```

ctg gat gat gga aga tta atg agt tcc ttt aat gat ctc atg caa cgc	446
Leu Asp Asp Gly Arg Leu Met Ser Ser Phe Asn Asp Leu Met Gln Arg	
115 120 125	
ctt tgg ata gtc gat agc gtt gaa cgt ttg ggg ata gct aga cat ttc	494
Leu Trp Ile Val Asp Ser Val Glu Arg Leu Gly Ile Ala Arg His Phe	
130 135 140	
aag aac gag ata aca tca gct ctg gat tat gtt ttc cgt tac tgg gag	542
Lys Asn Glu Ile Thr Ser Ala Leu Asp Tyr Val Phe Arg Tyr Trp Glu	
145 150 155	
gaa aac ggc att gga tgt ggg aga gac agt att gtt act gat ctc aac	590
Glu Asn Gly Ile Gly Cys Gly Arg Asp Ser Ile Val Thr Asp Leu Asn	
160 165 170	
tca act gcg ttg ggg ttt cga act ctt cga tta cac ggg tac act gta	638
Ser Thr Ala Leu Gly Phe Arg Thr Leu Arg Leu His Gly Tyr Thr Val	
175 180 185 190	
tct cca gag gtt tta aaa gct ttt caa gat caa aat gga cag ttt gta	686
Ser Pro Glu Val Leu Lys Ala Phe Gln Asp Gln Asn Gly Gln Phe Val	
195 200 205	
tgc tcc ccc ggt cag aca gag ggt gag atc aga agc gtt ctt aac tta	734
Cys Ser Pro Gly Gln Thr Glu Gly Glu Ile Arg Ser Val Leu Asn Leu	
210 215 220	
tat cgg gct tcc ctc att gcc ttc cct ggt gag aaa gtt atg gaa gaa	782
Tyr Arg Ala Ser Leu Ile Ala Phe Pro Gly Glu Lys Val Met Glu Glu	
225 230 235	
gct gaa atc ttc tcc aca aga tat ttg aaa gaa gct cta caa aag att	830
Ala Glu Ile Phe Ser Thr Arg Tyr Leu Lys Glu Ala Leu Gln Lys Ile	
240 245 250	
cca gtc tcc gct ctt tca caa gag ata aag ttt gtt atg gaa tat ggc	878
Pro Val Ser Ala Leu Ser Gln Glu Ile Lys Phe Val Met Glu Tyr Gly	
255 260 265 270	
tgg cac aca aat ttg cca aga ttg gaa gca aga aat tac ata gac aca	926
Trp His Thr Asn Leu Pro Arg Leu Glu Ala Arg Asn Tyr Ile Asp Thr	
275 280 285	
ctt gag aaa gac acc agt gca tgg ctc aat aaa aat gct ggg aag aag	974
Leu Glu Lys Asp Thr Ser Ala Trp Leu Asn Lys Asn Ala Gly Lys Lys	
290 295 300	
ctt tta gaa ctt gca aaa ttg gag ttc aat ata ttt aac tcc tta caa	1022
Leu Leu Glu Leu Ala Lys Leu Glu Phe Asn Ile Phe Asn Ser Leu Gln	
305 310 315	
caa aag gaa tta caa tat ctt ttg aga tgg tgg aaa gag tcg gat ttg	1070
Gln Lys Glu Leu Gln Tyr Leu Leu Arg Trp Trp Lys Glu Ser Asp Leu	
320 325 330	
cct aaa ttg aca ttt gct cgg cat cgt cat gtg gaa ttc tac act ttg	1118
Pro Lys Leu Thr Phe Ala Arg His Arg His Val Glu Phe Tyr Thr Leu	
335 340 345 350	

gcc tct tgt att gcc att gac cca aaa cat tct gca ttc aga cta ggc Ala Ser Cys Ile Ala Ile Asp Pro Lys His Ser Ala Phe Arg Leu Gly 355 360 365	1166
ttc gcc aaa atg tgt cat ctt gtc aca gtt ttg gac gat att tac gac Phe Ala Lys Met Cys His Leu Val Thr Val Leu Asp Asp Ile Tyr Asp 370 375 380	1214
act ttt gga acg att gac gag ctt gaa ctc ttc aca tct gca att aag Thr Phe Gly Thr Ile Asp Glu Leu Glu Leu Phe Thr Ser Ala Ile Lys 385 390 395	1262
aga tgg aat tca tca gag ata gaa cac ctt cca gaa tat atg aaa tgt Arg Trp Asn Ser Ser Glu Ile Glu His Leu Pro Glu Tyr Met Lys Cys 400 405 410	1310
gtg tac atg gtc gtg ttt gaa act gta aat gaa ctg aca cga gag gcg Val Tyr Met Val Val Phe Glu Thr Val Asn Glu Leu Thr Arg Glu Ala 415 420 425 430	1358
gag aag act caa ggg aga aac act ctc aac tat gtt cga aag gct tgg Glu Lys Thr Gln Gly Arg Asn Thr Leu Asn Tyr Val Arg Lys Ala Trp 435 440 445	1406
gag gct tat ttt gat tca tat atg gaa gaa gca aaa tgg atc tct aat Glu Ala Tyr Phe Asp Ser Tyr Met Glu Glu Ala Lys Trp Ile Ser Asn 450 455 460	1454
ggt tat ctg cca atg ttt gaa gag tac cat gag aat ggg aaa gtg agc Gly Tyr Leu Pro Met Phe Glu Glu Tyr His Glu Asn Gly Lys Val Ser 465 470 475	1502
tct gca tat cgc gta gca aca ttg caa ccc atc ctc act ttg aat gca Ser Ala Tyr Arg Val Ala Thr Leu Gln Pro Ile Leu Thr Leu Asn Ala 480 485 490	1550
tgg ctt cct gat tac atc ttg aag gga att gat ttt cca tcc agg ttc Trp Leu Pro Asp Tyr Ile Leu Lys Gly Ile Asp Phe Pro Ser Arg Phe 495 500 505 510	1598
aat gat ttg gca tcg tcc ttc ctt cgg cta cga ggt gac aca cgc tgc Asn Asp Leu Ala Ser Ser Phe Leu Arg Leu Arg Gly Asp Thr Arg Cys 515 520 525	1646
tac aag gcc gat agg gat cgt ggt gaa gaa gct tcg tgt ata tca tgt Tyr Lys Ala Asp Arg Asp Arg Gly Glu Glu Ala Ser Cys Ile Ser Cys 530 535 540	1694
tat atg aaa gac aat cct gga tca acc gaa gaa gat gcc ctc aat cat Tyr Met Lys Asp Asn Pro Gly Ser Thr Glu Glu Asp Ala Leu Asn His 545 550 555	1742
atc aat gcc atg gtc aat gac ata atc aaa gaa tta aat tgg gaa ctt Ile Asn Ala Met Val Asn Asp Ile Ile Lys Glu Leu Asn Trp Glu Leu 560 565 570	1790
cta aga tcc aac gac aat att cca atg ctg gcc aag aaa cat gct ttt Leu Arg Ser Asn Asp Asn Ile Pro Met Leu Ala Lys Lys His Ala Phe 575 580 585 590	1838

gac ata aca aga gct ctc cac cat ctc tac ata tat cga gat ggc ttt 1886
 Asp Ile Thr Arg Ala Leu His His Leu Tyr Ile Tyr Arg Asp Gly Phe
 595 600 605

agt gtt gcc aac aag gaa aca aaa aaa ttg gtt atg gaa aca ctc ctt 1934
 Ser Val Ala Asn Lys Glu Thr Lys Lys Leu Val Met Glu Thr Leu Leu
 610 615 620

gaa tct atg ctt ttt taa ctataaccat atccataata ataagctcat 1982
 Glu Ser Met Leu Phe
 625

aatgctaaat tattggcctt atgacatagt ttatgtatgt acttgtgtga attcaatcat 2042

atcgtgtggg tatgattaaa aagctagagc ttactagggt agtaacatgg tgataaaagt 2102

tataaaatgt gagttataga gatacccatg ttgaataatg aattacaaaa agagaaattt 2162

atgtagaata agattggaag cttttcaatt gttt 2196

<210> 2

<211> 627

<212> PRT

<213> *Abies grandis*

<400> 2

Met Ala Leu Val Ser Ile Ser Pro Leu Ala Ser Lys Ser Cys Leu Arg
 1 5 10 15

Lys Ser Leu Ile Ser Ser Ile His Glu His Lys Pro Pro Tyr Arg Thr
 20 25 30

Ile Pro Asn Leu Gly Met Arg Arg Arg Gly Lys Ser Val Thr Pro Ser
 35 40 45

Met Ser Ile Ser Leu Ala Thr Ala Ala Pro Asp Asp Gly Val Gln Arg
 50 55 60

Arg Ile Gly Asp Tyr His Ser Asn Ile Trp Asp Asp Asp Phe Ile Gln
 65 70 75 80

Ser Leu Ser Thr Pro Tyr Gly Glu Pro Ser Tyr Gln Glu Arg Ala Glu
 85 90 95

Arg Leu Ile Val Glu Val Lys Lys Ile Phe Asn Ser Met Tyr Leu Asp
 100 105 110

Asp Gly Arg Leu Met Ser Ser Phe Asn Asp Leu Met Gln Arg Leu Trp
 115 120 125

Ile Val Asp Ser Val Glu Arg Leu Gly Ile Ala Arg His Phe Lys Asn
 130 135 140

Glu Ile Thr Ser Ala Leu Asp Tyr Val Phe Arg Tyr Trp Glu Glu Asn
 145 150 155 160

Gly Ile Gly Cys Gly Arg Asp Ser Ile Val Thr Asp Leu Asn Ser Thr
 165 170 175

Ala Leu Gly Phe Arg Thr Leu Arg Leu His Gly Tyr Thr Val Ser Pro
 180 185 190
 Glu Val Leu Lys Ala Phe Gln Asp Gln Asn Gly Gln Phe Val Cys Ser
 195 200 205
 Pro Gly Gln Thr Glu Gly Glu Ile Arg Ser Val Leu Asn Leu Tyr Arg
 210 215 220
 Ala Ser Leu Ile Ala Phe Pro Gly Glu Lys Val Met Glu Glu Ala Glu
 225 230 235 240
 Ile Phe Ser Thr Arg Tyr Leu Lys Glu Ala Leu Gln Lys Ile Pro Val
 245 250 255
 Ser Ala Leu Ser Gln Glu Ile Lys Phe Val Met Glu Tyr Gly Trp His
 260 265 270
 Thr Asn Leu Pro Arg Leu Glu Ala Arg Asn Tyr Ile Asp Thr Leu Glu
 275 280 285
 Lys Asp Thr Ser Ala Trp Leu Asn Lys Asn Ala Gly Lys Lys Leu Leu
 290 295 300
 Glu Leu Ala Lys Leu Glu Phe Asn Ile Phe Asn Ser Leu Gln Gln Lys
 305 310 315 320
 Glu Leu Gln Tyr Leu Leu Arg Trp Trp Lys Glu Ser Asp Leu Pro Lys
 325 330 335
 Leu Thr Phe Ala Arg His Arg His Val Glu Phe Tyr Thr Leu Ala Ser
 340 345 350
 Cys Ile Ala Ile Asp Pro Lys His Ser Ala Phe Arg Leu Gly Phe Ala
 355 360 365
 Lys Met Cys His Leu Val Thr Val Leu Asp Asp Ile Tyr Asp Thr Phe
 370 375 380
 Gly Thr Ile Asp Glu Leu Glu Leu Phe Thr Ser Ala Ile Lys Arg Trp
 385 390 395 400
 Asn Ser Ser Glu Ile Glu His Leu Pro Glu Tyr Met Lys Cys Val Tyr
 405 410 415
 Met Val Val Phe Glu Thr Val Asn Glu Leu Thr Arg Glu Ala Glu Lys
 420 425 430
 Thr Gln Gly Arg Asn Thr Leu Asn Tyr Val Arg Lys Ala Trp Glu Ala
 435 440 445
 Tyr Phe Asp Ser Tyr Met Glu Glu Ala Lys Trp Ile Ser Asn Gly Tyr
 450 455 460
 Leu Pro Met Phe Glu Glu Tyr His Glu Asn Gly Lys Val Ser Ser Ala
 465 470 475 480
 Tyr Arg Val Ala Thr Leu Gln Pro Ile Leu Thr Leu Asn Ala Trp Leu
 485 490 495

Pro Asp Tyr Ile Leu Lys Gly Ile Asp Phe Pro Ser Arg Phe Asn Asp
500 505 510

Leu Ala Ser Ser Phe Leu Arg Leu Arg Gly Asp Thr Arg Cys Tyr Lys
515 520 525

Ala Asp Arg Asp Arg Gly Glu Glu Ala Ser Cys Ile Ser Cys Tyr Met
530 535 540

Lys Asp Asn Pro Gly Ser Thr Glu Glu Asp Ala Leu Asn His Ile Asn
545 550 555 560

Ala Met Val Asn Asp Ile Ile Lys Glu Leu Asn Trp Glu Leu Leu Arg
565 570 575

Ser Asn Asp Asn Ile Pro Met Leu Ala Lys Lys His Ala Phe Asp Ile
580 585 590

Thr Arg Ala Leu His His Leu Tyr Ile Tyr Arg Asp Gly Phe Ser Val
595 600 605

Ala Asn Lys Glu Thr Lys Lys Leu Val Met Glu Thr Leu Leu Glu Ser
610 615 620

Met Leu Phe
625

<210> 3

<211> 2018

<212> DNA

<213> *Abies grandis*

<220>

<221> CDS

<222> (6)..(1892)

<223> Clone AG3.18 encoding pinene synthase

<400> 3

cagca atg gct cta gtt tct acc gca ccg ttg gct tcc aaa tca tgc ctg 50
Met Ala Leu Val Ser Thr Ala Pro Leu Ala Ser Lys Ser Cys Leu
1 5 10 15

cac aaa tcg ttg atc agt tct acc cat gag ctt aag gct ctc tct aga 98
His Lys Ser Leu Ile Ser Ser Thr His Glu Leu Lys Ala Leu Ser Arg
20 25 30

aca att cca gct cta gga atg agt agg cga ggg aaa tct atc act cct 146
Thr Ile Pro Ala Leu Gly Met Ser Arg Arg Gly Lys Ser Ile Thr Pro
35 40 45

tcc atc agc atg agc tct acc acc gtt gta acc gat gat ggt gta cga 194
Ser Ile Ser Met Ser Ser Thr Thr Val Val Thr Asp Asp Gly Val Arg
50 55 60

aga cgc atg ggc gat ttc cat tcc aac ctc tgg gac gat gat gtc ata 242
Arg Arg Met Gly Asp Phe His Ser Asn Leu Trp Asp Asp Val Ile
65 70 75

cag tct tta cca acg gct tat gag gaa aaa tcg tac ctg gag cgt gct	290
Gln Ser Leu Pro Thr Ala Tyr Glu Glu Lys Ser Tyr Leu Glu Arg Ala	
80 85 90 95	
gag aaa ctg atc ggg gaa gta aag aac atg ttc aat tcg atg tca tta	338
Glu Lys Leu Ile Gly Glu Val Lys Asn Met Phe Asn Ser Met Ser Leu	
100 105 110	
gaa gat gga gag tta atg agt ccg ctc aat gat ctc att caa cgc ctt	386
Glu Asp Gly Glu Leu Met Ser Pro Leu Asn Asp Leu Ile Gln Arg Leu	
115 120 125	
tgg att gtc gac agc ctt gaa cgt ttg ggg atc cat aga cat ttc aaa	434
Trp Ile Val Asp Ser Leu Glu Arg Leu Gly Ile His Arg His Phe Lys	
130 135 140	
gat gag ata aaa tcg gcg ctt gat tat gtt tac agt tat tgg ggc gaa	482
Asp Glu Ile Lys Ser Ala Leu Asp Tyr Val Tyr Ser Tyr Trp Gly Glu	
145 150 155	
aat ggc atc gga tgc ggg agg gag agt gtt gtt act gat ctg aac tca	530
Asn Gly Ile Gly Cys Gly Arg Glu Ser Val Val Thr Asp Leu Asn Ser	
160 165 170 175	
act gcg ttg ggg ctt cga acc cta cga cta cac gga tac ccg gtg tct	578
Thr Ala Leu Gly Leu Arg Thr Leu Arg Leu His Gly Tyr Pro Val Ser	
180 185 190	
tca gat gtt ttc aaa gct ttc aaa ggc caa aat ggg cag ttt tcc tgc	626
Ser Asp Val Phe Lys Ala Phe Lys Gly Gln Asn Gly Gln Phe Ser Cys	
195 200 205	
tct gaa aat att cag aca gat gaa gag atc aga ggc gtt ctg aat tta	674
Ser Glu Asn Ile Gln Thr Asp Glu Glu Ile Arg Gly Val Leu Asn Leu	
210 215 220	
ttc cgg gcc tcc ctc att gcc ttt cca ggg gag aaa att atg gat gag	722
Phe Arg Ala Ser Leu Ile Ala Phe Pro Gly Glu Lys Ile Met Asp Glu	
225 230 235	
gct gaa atc ttc tct acc aaa tat tta aaa gaa gcc ctg caa aag att	770
Ala Glu Ile Phe Ser Thr Lys Tyr Leu Lys Glu Ala Leu Gln Lys Ile	
240 245 250 255	
ccg gtc tcc agt ctt tcg cga gag atc ggg gac gtt ttg gaa tat ggt	818
Pro Val Ser Ser Leu Ser Arg Glu Ile Gly Asp Val Leu Glu Tyr Gly	
260 265 270	
tgg cac aca tat ttg ccg cga ttg gaa gca agg aat tac atc caa gtc	866
Trp His Thr Tyr Leu Pro Arg Leu Glu Ala Arg Asn Tyr Ile Gln Val	
275 280 285	
ttt gga cag gac act gag aac acg aag tca tat gtg aag agc aaa aaa	914
Phe Gly Gln Asp Thr Glu Asn Thr Lys Ser Tyr Val Lys Ser Lys Lys	
290 295 300	
ctt tta gaa ctc gca aaa ttg gag ttc aac atc ttt caa tcc tta caa	962
Leu Leu Glu Leu Ala Lys Leu Glu Phe Asn Ile Phe Gln Ser Leu Gln	
305 310 315	

aag Lys 320	agg Arg	gag Glu	tta Leu	gaa Glu	agt Ser 325	ctg Leu	gtc Val	aga Arg	tgg Trp	tgg Trp 330	aaa Lys	gaa Glu	tcg Ser	ggt Gly	ttt Phe 335	1010
cct Pro	gag Glu	atg Met	acc Thr 340	ttc Phe	tgc Cys	cga Arg	cat His	cgt Arg	cac His 345	gtg Val	gaa Glu	tac Tyr	tac Tyr	act Thr 350	ttg Leu	1058
gct Ala	tcc Ser	tgc Cys	att Ile 355	gcg Ala	ttc Phe	gag Glu	cct Pro	caa Gln 360	cat His	tct Ser	gga Gly	ttc Phe	aga Arg 365	ctc Leu	ggc Gly	1106
ttt Phe	gcc Ala	aag Lys 370	acg Thr	tgt Cys	cat His	ctt Leu	atc Ile 375	acg Thr	gtt Val	ctt Leu	gac Asp	gat Asp 380	atg Met	tac Tyr	gac Asp	1154
acc Thr 385	ttc Phe	ggc Gly	aca Thr	gta Val	gac Asp	gag Glu 390	ctg Leu	gaa Glu	ctc Leu	ttc Phe	aca Thr 395	gcg Ala	aca Thr	atg Met	aag Lys	1202
aga Arg 400	tgg Trp	gat Asp	ccg Pro	tcc Ser	tcg Ser 405	ata Ile	gat Asp	tgc Cys	ctt Leu	cca Pro 410	gaa Glu	tat Tyr	atg Met	aaa Lys	gga Gly 415	1250
gtg Val	tac Tyr	ata Ile	gcg Ala	gtt Val	tac Tyr	gac Asp	acc Thr	gta Val	aat Asn 425	gaa Glu	atg Met	gct Ala	cga Arg	gag Glu 430	gca Ala	1298
gag Glu	gag Glu	gct Ala	caa Gln 435	ggc Gly	cga Arg	gat Asp	acg Thr	ctc Leu 440	aca Thr	tat Tyr	gct Ala	cgg Arg	gaa Glu 445	gct Ala	tgg Trp	1346
gag Glu	gct Ala	tat Tyr 450	att Ile	gat Asp	tcg Ser	tat Tyr	atg Met 455	caa Gln	gaa Glu	gca Ala	agg Arg	tgg Trp 460	atc Ile	gcc Ala	act Thr	1394
ggt Gly 465	tac Tyr	ctg Leu	ccc Pro	tcc Ser	ttt Phe	gat Asp 470	gag Glu	tac Tyr	tac Tyr	gag Glu	aat Asn 475	ggg Gly	aaa Lys	gtt Val	agc Ser	1442
tgt Cys 480	ggt Gly	cat His	cgc Arg	ata Ile	tcc Ser 485	gca Ala	ttg Leu	caa Gln	ccc Pro	att Ile 490	ctg Leu	aca Thr	atg Met	gac Asp	atc Ile 495	1490
ccc Pro	ttt Phe	cct Pro	gat Asp	cat His 500	atc Ile	ctc Leu	aag Lys	gaa Glu	gtt Val 505	gac Asp	ttc Phe	cca Pro	tca Ser	aag Lys 510	ctt Leu	1538
aac Asn	gac Asp	ttg Leu	gca Ala 515	tgt Cys	gcc Ala	atc Ile	ctt Leu	cga Arg 520	tta Leu	cga Arg	ggt Gly	gat Asp	acg Thr 525	cgg Arg	tgc Cys	1586
tac Tyr	aag Lys	gcg Ala 530	gac Asp	agg Arg	gct Ala	cgt Arg	gga Gly 535	gaa Glu	gaa Glu	gct Ala	tcc Ser	tct Ser	ata Ile	tca Ser	tgt Cys	1634
tat Tyr 545	atg Met	aaa Lys	gac Asp	aat Asn	cct Pro	gga Gly 550	gta Val	tca Ser	gag Glu	gaa Glu	gat Asp 555	gct Ala	ctc Leu	gat Asp	cat His	1682

atc aac gcc atg atc agt gac gta atc aaa gga tta aat tgg gaa ctt 1730
 Ile Asn Ala Met Ile Ser Asp Val Ile Lys Gly Leu Asn Trp Glu Leu
 560 565 570 575

ctc aaa cca gac atc aat gtt ccc atc tcg gcg aag aaa cat gct ttt 1778
 Leu Lys Pro Asp Ile Asn Val Pro Ile Ser Ala Lys Lys His Ala Phe
 580 585 590

gac atc gcc aga gct ttc cat tac ggc tac aaa tac cga gac ggc tac 1826
 Asp Ile Ala Arg Ala Phe His Tyr Gly Tyr Lys Tyr Arg Asp Gly Tyr
 595 600 605

agc gtt gcc aac gtt gaa acg aag agt ttg gtc acg aga acc ctc ctt 1874
 Ser Val Ala Asn Val Glu Thr Lys Ser Leu Val Thr Arg Thr Leu Leu
 610 615 620

gaa tct gtg cct ttg tag caacagctca aatctatgcc ctatgctatg 1922
 Glu Ser Val Pro Leu
 625

tcgggttaaa atatatgtgg aaggtagccg ttggatgtag aggataagtt tgttataatt 1982
 taataaagtt gtaatttaaa aaaaaaaaaa aaaaaa 2018

<210> 4
 <211> 628
 <212> PRT
 <213> Abies grandis

<400> 4
 Met Ala Leu Val Ser Thr Ala Pro Leu Ala Ser Lys Ser Cys Leu His
 1 5 10 15
 Lys Ser Leu Ile Ser Ser Thr His Glu Leu Lys Ala Leu Ser Arg Thr
 20 25 30
 Ile Pro Ala Leu Gly Met Ser Arg Arg Gly Lys Ser Ile Thr Pro Ser
 35 40 45
 Ile Ser Met Ser Ser Thr Thr Val Val Thr Asp Asp Gly Val Arg Arg
 50 55 60
 Arg Met Gly Asp Phe His Ser Asn Leu Trp Asp Asp Asp Val Ile Gln
 65 70 75 80
 Ser Leu Pro Thr Ala Tyr Glu Glu Lys Ser Tyr Leu Glu Arg Ala Glu
 85 90 95
 Lys Leu Ile Gly Glu Val Lys Asn Met Phe Asn Ser Met Ser Leu Glu
 100 105 110
 Asp Gly Glu Leu Met Ser Pro Leu Asn Asp Leu Ile Gln Arg Leu Trp
 115 120 125
 Ile Val Asp Ser Leu Glu Arg Leu Gly Ile His Arg His Phe Lys Asp
 130 135 140
 Glu Ile Lys Ser Ala Leu Asp Tyr Val Tyr Ser Tyr Trp Gly Glu Asn
 145 150 155 160

Gly Ile Gly Cys Gly Arg Glu Ser Val Val Thr Asp Leu Asn Ser Thr
 165 170 175
 Ala Leu Gly Leu Arg Thr Leu Arg Leu His Gly Tyr Pro Val Ser Ser
 180 185 190
 Asp Val Phe Lys Ala Phe Lys Gly Gln Asn Gly Gln Phe Ser Cys Ser
 195 200 205
 Glu Asn Ile Gln Thr Asp Glu Glu Ile Arg Gly Val Leu Asn Leu Phe
 210 215 220
 Arg Ala Ser Leu Ile Ala Phe Pro Gly Glu Lys Ile Met Asp Glu Ala
 225 230 235 240
 Glu Ile Phe Ser Thr Lys Tyr Leu Lys Glu Ala Leu Gln Lys Ile Pro
 245 250 255
 Val Ser Ser Leu Ser Arg Glu Ile Gly Asp Val Leu Glu Tyr Gly Trp
 260 265 270
 His Thr Tyr Leu Pro Arg Leu Glu Ala Arg Asn Tyr Ile Gln Val Phe
 275 280 285
 Gly Gln Asp Thr Glu Asn Thr Lys Ser Tyr Val Lys Ser Lys Lys Leu
 290 295 300
 Leu Glu Leu Ala Lys Leu Glu Phe Asn Ile Phe Gln Ser Leu Gln Lys
 305 310 315 320
 Arg Glu Leu Glu Ser Leu Val Arg Trp Trp Lys Glu Ser Gly Phe Pro
 325 330 335
 Glu Met Thr Phe Cys Arg His Arg His Val Glu Tyr Tyr Thr Leu Ala
 340 345 350
 Ser Cys Ile Ala Phe Glu Pro Gln His Ser Gly Phe Arg Leu Gly Phe
 355 360 365
 Ala Lys Thr Cys His Leu Ile Thr Val Leu Asp Asp Met Tyr Asp Thr
 370 375 380
 Phe Gly Thr Val Asp Glu Leu Glu Leu Phe Thr Ala Thr Met Lys Arg
 385 390 395 400
 Trp Asp Pro Ser Ser Ile Asp Cys Leu Pro Glu Tyr Met Lys Gly Val
 405 410 415
 Tyr Ile Ala Val Tyr Asp Thr Val Asn Glu Met Ala Arg Glu Ala Glu
 420 425 430
 Glu Ala Gln Gly Arg Asp Thr Leu Thr Tyr Ala Arg Glu Ala Trp Glu
 435 440 445
 Ala Tyr Ile Asp Ser Tyr Met Gln Glu Ala Arg Trp Ile Ala Thr Gly
 450 455 460
 Tyr Leu Pro Ser Phe Asp Glu Tyr Tyr Glu Asn Gly Lys Val Ser Cys
 465 470 475 480

Gly His Arg Ile Ser Ala Leu Gln Pro Ile Leu Thr Met Asp Ile Pro
 485 490 495
 Phe Pro Asp His Ile Leu Lys Glu Val Asp Phe Pro Ser Lys Leu Asn
 500 505 510
 Asp Leu Ala Cys Ala Ile Leu Arg Leu Arg Gly Asp Thr Arg Cys Tyr
 515 520 525
 Lys Ala Asp Arg Ala Arg Gly Glu Glu Ala Ser Ser Ile Ser Cys Tyr
 530 535 540
 Met Lys Asp Asn Pro Gly Val Ser Glu Glu Asp Ala Leu Asp His Ile
 545 550 555 560
 Asn Ala Met Ile Ser Asp Val Ile Lys Gly Leu Asn Trp Glu Leu Leu
 565 570 575
 Lys Pro Asp Ile Asn Val Pro Ile Ser Ala Lys Lys His Ala Phe Asp
 580 585 590
 Ile Ala Arg Ala Phe His Tyr Gly Tyr Lys Tyr Arg Asp Gly Tyr Ser
 595 600 605
 Val Ala Asn Val Glu Thr Lys Ser Leu Val Thr Arg Thr Leu Leu Glu
 610 615 620
 Ser Val Pro Leu
 625

<210> 5
 <211> 2089
 <212> DNA
 <213> Abies grandis

 <220>
 <221> CDS
 <222> (73)..(1986)
 <223> Clone AG10 encoding limonene synthase

<400> 5
 tgccgttttaa tcggttttaaa gaagctacca tagttcgggt taaagaagct accatagttt 60
 aggcaggaat cc atg gct ctc ctt tct atc gta tct ttg cag gtt ccc aaa 111
 Met Ala Leu Leu Ser Ile Val Ser Leu Gln Val Pro Lys
 1 5 10
 tcc tgc ggg ctg aaa tcg ttg atc agt tcc agc aat gtg cag aag gct 159
 Ser Cys Gly Leu Lys Ser Leu Ile Ser Ser Ser Asn Val Gln Lys Ala
 15 20 25
 ctc tgt atc tct aca gca gtc cca aca ctc aga atg cgt agg cga cag 207
 Leu Cys Ile Ser Thr Ala Val Pro Thr Leu Arg Met Arg Arg Arg Gln
 30 35 40 45
 aaa gct ctg gtc atc aac atg aaa ttg acc act gta tcc cat cgt gat 255
 Lys Ala Leu Val Ile Asn Met Lys Leu Thr Thr Val Ser His Arg Asp
 50 55 60

gat aat ggt ggt ggt gta ctg caa aga cgc ata gcc gat cat cat ccc 303
 Asp Asn Gly Gly Gly Val Leu Gln Arg Arg Ile Ala Asp His His Pro
 65 70 75

aac ctg tgg gaa gat gat ttc ata caa tca ttg tcc tca cct tat ggg 351
 Asn Leu Trp Glu Asp Asp Phe Ile Gln Ser Leu Ser Ser Pro Tyr Gly
 80 85 90

gga tct tcg tac agt gaa cgt gct gag aca gtc gtt gag gaa gta aaa 399
 Gly Ser Ser Tyr Ser Glu Arg Ala Glu Thr Val Val Glu Glu Val Lys
 95 100 105

gag atg ttc aat tca ata cca aat aat aga gaa tta ttt ggt tcc caa 447
 Glu Met Phe Asn Ser Ile Pro Asn Asn Arg Glu Leu Phe Gly Ser Gln
 110 115 120 125

aat gat ctc ctt aca cgc ctt tgg atg gtg gat agc att gaa cgt ctg 495
 Asn Asp Leu Leu Thr Arg Leu Trp Met Val Asp Ser Ile Glu Arg Leu
 130 135 140

ggg ata gat aga cat ttc caa aat gag ata aga gta gcc ctc gat tat 543
 Gly Ile Asp Arg His Phe Gln Asn Glu Ile Arg Val Ala Leu Asp Tyr
 145 150 155

gtt tac agt tat tgg aag gaa aag gaa ggc att ggg tgt ggc aga gat 591
 Val Tyr Ser Tyr Trp Lys Glu Lys Glu Gly Ile Gly Cys Gly Arg Asp
 160 165 170

tct act ttt cct gat ctc aac tcg act gcc ttg gcg ctt cga act ctt 639
 Ser Thr Phe Pro Asp Leu Asn Ser Thr Ala Leu Ala Leu Arg Thr Leu
 175 180 185

cga ctg cac gga tac aat gtg tct tca gat gtg ctg gaa tac ttc aaa 687
 Arg Leu His Gly Tyr Asn Val Ser Ser Asp Val Leu Glu Tyr Phe Lys
 190 195 200 205

gat gaa aag ggg cat ttt gcc tgc cct gca atc cta acc gag gga cag 735
 Asp Glu Lys Gly His Phe Ala Cys Pro Ala Ile Leu Thr Glu Gly Gln
 210 215 220

atc act aga agt gtt cta aat tta tat cgg gct tcc ctg gtc gcc ttt 783
 Ile Thr Arg Ser Val Leu Asn Leu Tyr Arg Ala Ser Leu Val Ala Phe
 225 230 235

ccc ggg gag aaa gtt atg gaa gag gct gaa atc ttc tcg gca tct tat 831
 Pro Gly Glu Lys Val Met Glu Glu Ala Glu Ile Phe Ser Ala Ser Tyr
 240 245 250

ttg aaa aaa gtc tta caa aag att ccg gtc tcc aat ctt tca gga gag 879
 Leu Lys Lys Val Leu Gln Lys Ile Pro Val Ser Asn Leu Ser Gly Glu
 255 260 265

ata gaa tat gtt ttg gaa tat ggt tgg cac acg aat ttg ccg aga ttg 927
 Ile Glu Tyr Val Leu Glu Tyr Gly Trp His Thr Asn Leu Pro Arg Leu
 270 275 280 285

gaa gca aga aat tat atc gag gtc tac gag cag agc ggc tat gaa agc 975
 Glu Ala Arg Asn Tyr Ile Glu Val Tyr Glu Gln Ser Gly Tyr Glu Ser
 290 295 300

tta aac gag atg cca tat atg aac atg aag aag ctt tta caa ctt gca Leu Asn Glu Met Pro Tyr Met Asn Met Lys Lys Leu Leu Gln Leu Ala	1023
305 310 315	
aaa ttg gag ttc aat atc ttt cac tct ttg caa cta aga gag tta caa Lys Leu Glu Phe Asn Ile Phe His Ser Leu Gln Leu Arg Glu Leu Gln	1071
320 325 330	
tct atc tcc aga tgg tgg aaa gaa tca ggt tgc tct caa ctg act ttt Ser Ile Ser Arg Trp Trp Lys Glu Ser Gly Ser Ser Gln Leu Thr Phe	1119
335 340 345	
aca cgg cat cgt cac gtg gaa tac tac act atg gca tct tgc att tct Thr Arg His Arg His Val Glu Tyr Tyr Thr Met Ala Ser Cys Ile Ser	1167
350 355 360 365	
atg ttg cca aaa cat tca gct ttc aga atg gag ttt gtc aaa gtg tgt Met Leu Pro Lys His Ser Ala Phe Arg Met Glu Phe Val Lys Val Cys	1215
370 375 380	
cat ctt gta aca gtt ctc gat gat ata tat gac act ttt gga aca atg His Leu Val Thr Val Leu Asp Asp Ile Tyr Asp Thr Phe Gly Thr Met	1263
385 390 395	
aac gaa ctc caa ctt ttt acg gat gca att aag aga tgg gat ttg tca Asn Glu Leu Gln Leu Phe Thr Asp Ala Ile Lys Arg Trp Asp Leu Ser	1311
400 405 410	
acg aca agg tgg ctt cca gaa tat atg aaa gga gtg tac atg gac ttg Thr Thr Arg Trp Leu Pro Glu Tyr Met Lys Gly Val Tyr Met Asp Leu	1359
415 420 425	
tat caa tgc att aat gaa atg gtg gaa gag gct gag aag act caa ggc Tyr Gln Cys Ile Asn Glu Met Val Glu Glu Ala Glu Lys Thr Gln Gly	1407
430 435 440 445	
cga gat atg ctc aac tat att caa aat gct tgg gaa gcc cta ttt gat Arg Asp Met Leu Asn Tyr Ile Gln Asn Ala Trp Glu Ala Leu Phe Asp	1455
450 455 460	
acc ttt atg caa gaa gca aag tgg atc tcc agc agt tat ctc cca acg Thr Phe Met Gln Glu Ala Lys Trp Ile Ser Ser Ser Tyr Leu Pro Thr	1503
465 470 475	
ttt gag gag tac ttg aag aat gca aaa gtt agt tct ggt tct cgc ata Phe Glu Glu Tyr Leu Lys Asn Ala Lys Val Ser Ser Gly Ser Arg Ile	1551
480 485 490	
gcc aca tta caa ccc att ctc act ttg gat gta cca ctt cct gat tac Ala Thr Leu Gln Pro Ile Leu Thr Leu Asp Val Pro Leu Pro Asp Tyr	1599
495 500 505	
ata ctg caa gaa att gat tat cca tcc aga ttc aat gag tta gct tgc Ile Leu Gln Glu Ile Asp Tyr Pro Ser Arg Phe Asn Glu Leu Ala Ser	1647
510 515 520 525	
tcc atc ctt cga cta cga ggt gac acg cgc tgc tac aag gcg gat agg Ser Ile Leu Arg Leu Arg Gly Asp Thr Arg Cys Tyr Lys Ala Asp Arg	1695
530 535 540	

gcc cgt gga gaa gaa gct tca gct ata tcg tgt tat atg aaa gac cat 1743
 Ala Arg Gly Glu Glu Ala Ser Ala Ile Ser Cys Tyr Met Lys Asp His
 545 550 555

cct gga tca ata gag gaa gat gct ctc aat cat atc aac gcc atg atc 1791
 Pro Gly Ser Ile Glu Glu Asp Ala Leu Asn His Ile Asn Ala Met Ile
 560 565 570

agt gat gca atc aga gaa tta aat tgg gag ctt ctc aga ccg gat agc 1839
 Ser Asp Ala Ile Arg Glu Leu Asn Trp Glu Leu Leu Arg Pro Asp Ser
 575 580 585

aaa agt ccc atc tct tcc aag aaa cat gct ttt gac atc acc aga gct 1887
 Lys Ser Pro Ile Ser Ser Lys Lys His Ala Phe Asp Ile Thr Arg Ala
 590 595 600 605

ttc cat cat gtc tac aaa tat cga gat ggt tac act gtt tcc aac aac 1935
 Phe His His Val Tyr Lys Tyr Arg Asp Gly Tyr Thr Val Ser Asn Asn
 610 615 620

gaa aca aag aat ttg gtg atg aaa acc gtt ctt gaa cct ctc gct ttg 1983
 Glu Thr Lys Asn Leu Val Met Lys Thr Val Leu Glu Pro Leu Ala Leu
 625 630 635

taa aaacatatag aatgcattaa aatgtgggaa gtctataatc tagactattc 2036

tctatctttc ataatgtaga tctggatgtg tattgaactc taaaaaaaaa aaa 2089

<210> 6
 <211> 637
 <212> PRT
 <213> *Abies grandis*

<400> 6
 Met Ala Leu Leu Ser Ile Val Ser Leu Gln Val Pro Lys Ser Cys Gly
 1 5 10 15
 Leu Lys Ser Leu Ile Ser Ser Ser Asn Val Gln Lys Ala Leu Cys Ile
 20 25 30
 Ser Thr Ala Val Pro Thr Leu Arg Met Arg Arg Arg Gln Lys Ala Leu
 35 40 45
 Val Ile Asn Met Lys Leu Thr Thr Val Ser His Arg Asp Asp Asn Gly
 50 55 60
 Gly Gly Val Leu Gln Arg Arg Ile Ala Asp His His Pro Asn Leu Trp
 65 70 75 80
 Glu Asp Asp Phe Ile Gln Ser Leu Ser Ser Pro Tyr Gly Gly Ser Ser
 85 90 95
 Tyr Ser Glu Arg Ala Glu Thr Val Val Glu Glu Val Lys Glu Met Phe
 100 105 110
 Asn Ser Ile Pro Asn Asn Arg Glu Leu Phe Gly Ser Gln Asn Asp Leu
 115 120 125

Leu Thr Arg Leu Trp Met Val Asp Ser Ile Glu Arg Leu Gly Ile Asp
 130 135 140
 Arg His Phe Gln Asn Glu Ile Arg Val Ala Leu Asp Tyr Val Tyr Ser
 145 150 155 160
 Tyr Trp Lys Glu Lys Glu Gly Ile Gly Cys Gly Arg Asp Ser Thr Phe
 165 170 175
 Pro Asp Leu Asn Ser Thr Ala Leu Ala Leu Arg Thr Leu Arg Leu His
 180 185 190
 Gly Tyr Asn Val Ser Ser Asp Val Leu Glu Tyf Phe Lys Asp Glu Lys
 195 200 205
 Gly His Phe Ala Cys Pro Ala Ile Leu Thr Glu Gly Gln Ile Thr Arg
 210 215 220
 Ser Val Leu Asn Leu Tyr Arg Ala Ser Leu Val Ala Phe Pro Gly Glu
 225 230 235 240
 Lys Val Met Glu Glu Ala Glu Ile Phe Ser Ala Ser Tyr Leu Lys Lys
 245 250 255
 Val Leu Gln Lys Ile Pro Val Ser Asn Leu Ser Gly Glu Ile Glu Tyr
 260 265 270
 Val Leu Glu Tyr Gly Trp His Thr Asn Leu Pro Arg Leu Glu Ala Arg
 275 280 285
 Asn Tyr Ile Glu Val Tyr Glu Gln Ser Gly Tyr Glu Ser Leu Asn Glu
 290 295 300
 Met Pro Tyr Met Asn Met Lys Lys Leu Leu Gln Leu Ala Lys Leu Glu
 305 310 315 320
 Phe Asn Ile Phe His Ser Leu Gln Leu Arg Glu Leu Gln Ser Ile Ser
 325 330 335
 Arg Trp Trp Lys Glu Ser Gly Ser Ser Gln Leu Thr Phe Thr Arg His
 340 345 350
 Arg His Val Glu Tyr Tyr Thr Met Ala Ser Cys Ile Ser Met Leu Pro
 355 360 365
 Lys His Ser Ala Phe Arg Met Glu Phe Val Lys Val Cys His Leu Val
 370 375 380
 Thr Val Leu Asp Asp Ile Tyr Asp Thr Phe Gly Thr Met Asn Glu Leu
 385 390 395 400
 Gln Leu Phe Thr Asp Ala Ile Lys Arg Trp Asp Leu Ser Thr Thr Arg
 405 410 415
 Trp Leu Pro Glu Tyr Met Lys Gly Val Tyr Met Asp Leu Tyr Gln Cys
 420 425 430
 Ile Asn Glu Met Val Glu Glu Ala Glu Lys Thr Gln Gly Arg Asp Met
 435 440 445

Leu Asn Tyr Ile Gln Asn Ala Trp Glu Ala Leu Phe Asp Thr Phe Met
 450 455 460
 Gln Glu Ala Lys Trp Ile Ser Ser Ser Tyr Leu Pro Thr Phe Glu Glu
 465 470 475 480
 Tyr Leu Lys Asn Ala Lys Val Ser Ser Gly Ser Arg Ile Ala Thr Leu
 485 490 495
 Gln Pro Ile Leu Thr Leu Asp Val Pro Leu Pro Asp Tyr Ile Leu Gln
 500 505 510
 Glu Ile Asp Tyr Pro Ser Arg Phe Asn Glu Leu Ala Ser Ser Ile Leu
 515 520 525
 Arg Leu Arg Gly Asp Thr Arg Cys Tyr Lys Ala Asp Arg Ala Arg Gly
 530 535 540
 Glu Glu Ala Ser Ala Ile Ser Cys Tyr Met Lys Asp His Pro Gly Ser
 545 550 555 560
 Ile Glu Glu Asp Ala Leu Asn His Ile Asn Ala Met Ile Ser Asp Ala
 565 570 575
 Ile Arg Glu Leu Asn Trp Glu Leu Leu Arg Pro Asp Ser Lys Ser Pro
 580 585 590
 Ile Ser Ser Lys Lys His Ala Phe Asp Ile Thr Arg Ala Phe His His
 595 600 605
 Val Tyr Lys Tyr Arg Asp Gly Tyr Thr Val Ser Asn Asn Glu Thr Lys
 610 615 620
 Asn Leu Val Met Lys Thr Val Leu Glu Pro Leu Ala Leu
 625 630 635

<210> 7

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Degenerate oligonucleotide PCR primer A wherein the letter "n" indicates an inosine residue

<220>

<221> misc_feature

<222> (1)..(25)

<223> Degenerate oligonucleotide Primer A wherein n represents inosine

<400> 7

arraygarra nggnrartay aarga

25

<210> 8

<211> 20

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: degenerate
oligonucleotide PCR primer B wherein the letter
"n" represents an inosine residue

<220>
<221> misc_feature
<222> (1)..(20)
<223> oligonucleotide PCR primer B wherein the letter n
represents an inosine residue

<400> 8
atgytncary tntaygargc

20

<210> 9
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: degenerate
oligonucleotide PCR primer C wherein the letter
"n" represents an inosine residue

<220>
<221> misc_feature
<222> (1)..(24)
<223> PCR primer C wherein the letter n represents
inosine

<400> 9
ctnkynrang gnctratrta ckty

24

<210> 10
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: degenerate
oligonucleotide PCR primer D wherein the letter
"n" represents an inosine residue

<220>
<221> misc_feature
<222> (1)..(23)
<223> PCR primer D wherein the letter n represents
inosine

<400> 10
gaygaynnnt wygaygcnya ygg

23

<210> 11
<211> 108

<212> DNA

<213> Artificial Sequence

<400> 11

gatgatgggt ttgatgcgca cggaacccta gatgaattga agctattcac tgaggctgtg 60
 agaagatggg acctctcctt tacagacaac ttccccgatt acatgaaa 108

<210> 12

<211> 104

<212> DNA

<213> *Abies grandis*

<400> 12

gacgacgggt atgatgcgca tggaacgatt gacgagcttg aactcttcac atctgcaatt 60
 aagagatgga attcatcaga gatagacagc ttccccgact atat 104

<210> 13

<211> 105

<212> DNA

<213> *Abies grandis*

<400> 13

gatgatgggt atgatgcgta cggaacgcttg gaagaaatca aaatcatgac agaggggagtg 60
 agacgatggg atctttcggt gaccgcttnc cccgactata tgaaa 105

<210> 14

<211> 117

<212> DNA

<213> *Abies grandis*

<400> 14

gacgatgggt atgatgcgca tggaaccttg gaccaactca aaatctttac agaggggagtg 60
 agacgatggg atgtttcggt ggtagaccac ttncctcgac tacatgcaat ctagacc 117

<210> 15

<211> 2424

<212> DNA

<213> *Abies grandis*

<220>

<221> CDS

<222> (2)..(2350)

<223> Clone AG1.28

<400> 15

g ggt tat gat ctt gtg cat tct ctt aaa tca cct tat att gat tct agt 49
 Gly Tyr Asp Leu Val His Ser Leu Lys Ser Pro Tyr Ile Asp Ser Ser
 1 5 10 15

tac aga gaa cgc gcg gag gtc ctt gtt agc gag att aaa gtg atg ctt 97
 Tyr Arg Glu Arg Ala Glu Val Leu Val Ser Glu Ile Lys Val Met Leu
 20 25 30

aat cca gct att aca gga gat gga gaa tca atg att act cca tct gct	145
Asn Pro Ala Ile Thr Gly Asp Gly Glu Ser Met Ile Thr Pro Ser Ala	
35 40 45	
tat gac aca gca tgg gta gcg agg gtg ccc gcc att gat ggc tct gct	193
Tyr Asp Thr Ala Trp Val Ala Arg Val Pro Ala Ile Asp Gly Ser Ala	
50 55 60	
cgc ccg caa ttt ccc caa aca gtt gac tgg att ttg aaa aac cag tta	241
Arg Pro Gln Phe Pro Gln Thr Val Asp Trp Ile Leu Lys Asn Gln Leu	
65 70 75 80	
aaa gat ggt tca tgg gga att cag tcc cac ttt ctg ctg tcc gac cgt	289
Lys Asp Gly Ser Trp Gly Ile Gln Ser His Phe Leu Leu Ser Asp Arg	
85 90 95	
ctt ctt gcc act ctt tct tgt gtt ctt gtg ctc ctt aaa tgg aac gtt	337
Leu Leu Ala Thr Leu Ser Cys Val Leu Val Leu Leu Lys Trp Asn Val	
100 105 110	
ggg gat ctg caa gta gag cag gga att gaa ttc ata aag agc aat ctg	385
Gly Asp Leu Gln Val Glu Gln Gly Ile Glu Phe Ile Lys Ser Asn Leu	
115 120 125	
gaa cta gta aag gat gaa acc gat caa gat agc ttg gta aca gac ttt	433
Glu Leu Val Lys Asp Glu Thr Asp Gln Asp Ser Leu Val Thr Asp Phe	
130 135 140	
gag atc ata ttt cct tct ctg tta aga gaa gct caa tct ctg cgc ctc	481
Glu Ile Ile Phe Pro Ser Leu Leu Arg Glu Ala Gln Ser Leu Arg Leu	
145 150 155 160	
gga ctt ccc tac gac ctg cct tat ata cat ctg ttg cag act aaa cgg	529
Gly Leu Pro Tyr Asp Leu Pro Tyr Ile His Leu Leu Gln Thr Lys Arg	
165 170 175	
cag gaa aga tta gca aaa ctt tca agg gag gaa att tat gcg gtt ccg	577
Gln Glu Arg Leu Ala Lys Leu Ser Arg Glu Glu Ile Tyr Ala Val Pro	
180 185 190	
tcg cca ttg ttg tat tct tta gag gga ata caa gat ata gtt gaa tgg	625
Ser Pro Leu Leu Tyr Ser Leu Glu Gly Ile Gln Asp Ile Val Glu Trp	
195 200 205	
gaa cga ata atg gaa gtt caa agt cag gat ggg tct ttc tta agc tca	673
Glu Arg Ile Met Glu Val Gln Ser Gln Asp Gly Ser Phe Leu Ser Ser	
210 215 220	
cct gct tct act gcc tgc gtt ttc atg cac aca gga gac gcg aaa tgc	721
Pro Ala Ser Thr Ala Cys Val Phe Met His Thr Gly Asp Ala Lys Cys	
225 230 235 240	
ctt gaa ttc ttg aac agt gtg atg atc aag ttt gga aat ttt gtt ccc	769
Leu Glu Phe Leu Asn Ser Val Met Ile Lys Phe Gly Asn Phe Val Pro	
245 250 255	
tgc ctg tat cct gtg gat ctg ctg gaa cgc ctg ttg atc gta gat aat	817
Cys Leu Tyr Pro Val Asp Leu Leu Glu Arg Leu Leu Ile Val Asp Asn	
260 265 270	

att gta cgc ctt gga atc tat aga cac ttt gaa aag gaa atc aag gaa Ile Val Arg Leu Gly Ile Tyr Arg His Phe Glu Lys Glu Ile Lys Glu 275 280 285	865
gct ctt gat tat gtt tac agg cat tgg aac gaa aga gga att ggg tgg Ala Leu Asp Tyr Val Tyr Arg His Trp Asn Glu Arg Gly Ile Gly Trp 290 295 300	913
ggc aga cta aat ccc ata gca gat ctt gag acc act gct ttg gga ttt Gly Arg Leu Asn Pro Ile Ala Asp Leu Glu Thr Thr Ala Leu Gly Phe 305 310 315 320	961
cga ttg ctt cgg ctg cat agg tac aat gta tct cca gcc att ttt gac Arg Leu Leu Arg Leu His Arg Tyr Asn Val Ser Pro Ala Ile Phe Asp 325 330 335	1009
aac ttc aaa gat gcc aat ggg aaa ttc att tgc tcg acc ggt caa ttc Asn Phe Lys Asp Ala Asn Gly Lys Phe Ile Cys Ser Thr Gly Gln Phe 340 345 350	1057
aac aaa gat gta gca agc atg ctg aat ctt tat aga gct tcc cag ctc Asn Lys Asp Val Ala Ser Met Leu Asn Leu Tyr Arg Ala Ser Gln Leu 355 360 365	1105
gca ttt ccc gga gaa aac att ctt gat gaa gct aaa agc ttc gct act Ala Phe Pro Gly Glu Asn Ile Leu Asp Glu Ala Lys Ser Phe Ala Thr 370 375 380	1153
aaa tat ttg aga gaa gct ctt gag aaa agt gag act tcc agt gca tgg Lys Tyr Leu Arg Glu Ala Leu Glu Lys Ser Glu Thr Ser Ser Ala Trp 385 390 395 400	1201
aac aac aaa caa aac ctg agc caa gag atc aaa tac gcg ctg aag act Asn Asn Lys Gln Asn Leu Ser Gln Glu Ile Lys Tyr Ala Leu Lys Thr 405 410 415	1249
tct tgg cat gcc agt gtt ccg aga gtg gaa gca aag aga tac tgt caa Ser Trp His Ala Ser Val Pro Arg Val Glu Ala Lys Arg Tyr Cys Gln 420 425 430	1297
gtg tat cgc cca gat tat gca cgc ata gca aaa tgc gtt tac aag cta Val Tyr Arg Pro Asp Tyr Ala Arg Ile Ala Lys Cys Val Tyr Lys Leu 435 440 445	1345
ccc tac gtg aac aat gaa aag ttt tta gag ctg gga aaa tta gat ttc Pro Tyr Val Asn Asn Glu Lys Phe Leu Glu Leu Gly Lys Leu Asp Phe 450 455 460	1393
aac att atc cag tcc atc cac caa gaa gaa atg aag aat gtt acc agc Asn Ile Ile Gln Ser Ile His Gln Glu Glu Met Lys Asn Val Thr Ser 465 470 475 480	1441
tgg ttt aga gat tcg ggg ttg cca cta ttc acc ttc gct cgg gag agg Trp Phe Arg Asp Ser Gly Leu Pro Leu Phe Thr Phe Ala Arg Glu Arg 485 490 495	1489
ccg ctg gaa ttc tac ttc tta gta gcg gcg ggg acc tat gaa ccc cag Pro Leu Glu Phe Tyr Phe Leu Val Ala Ala Gly Thr Tyr Glu Pro Gln 500 505 510	1537

tat gcc aaa tgc agg ttc ctc ttt aca aaa gtg gca tgc ttg cag act	1585
Tyr Ala Lys Cys Arg Phe Leu Phe Thr Lys Val Ala Cys Leu Gln Thr	
515 520 525	
gtt ctg gac gat atg tat gac act tat gga acc cta gat gaa ttg aag	1633
Val Leu Asp Asp Met Tyr Asp Thr Tyr Gly Thr Leu Asp Glu Leu Lys	
530 535 540	
cta ttc act gag gct gtg aga aga tgg gac ctc tcc ttt aca gaa aac	1681
Leu Phe Thr Glu Ala Val Arg Arg Trp Asp Leu Ser Phe Thr Glu Asn	
545 550 555 560	
ctt cca gac tat atg aaa cta tgt tac caa atc tat tat gac ata gtt	1729
Leu Pro Asp Tyr Met Lys Leu Cys Tyr Gln Ile Tyr Tyr Asp Ile Val	
565 570 575	
cac gag gtg gct tgg gag gca gag aag gaa cag ggg cgt gaa ttg gtc	1777
His Glu Val Ala Trp Glu Ala Glu Lys Glu Gln Gly Arg Glu Leu Val	
580 585 590	
agc ttt ttc aga aag gga tgg gag gat tat ctt ctg ggt tat tat gaa	1825
Ser Phe Phe Arg Lys Gly Trp Glu Asp Tyr Leu Leu Gly Tyr Tyr Glu	
595 600 605	
gaa gct gaa tgg tta gct gct gag tat gtg cct acc ttg gac gag tac	1873
Glu Ala Glu Trp Leu Ala Ala Glu Tyr Val Pro Thr Leu Asp Glu Tyr	
610 615 620	
ata aag aat gga atc aca tct atc ggc caa cgt ata ctt ctg ttg agt	1921
Ile Lys Asn Gly Ile Thr Ser Ile Gly Gln Arg Ile Leu Leu Leu Ser	
625 630 635 640	
gga gtg ttg ata atg gat ggg caa ctc ctt tcg caa gag gca tta gag	1969
Gly Val Leu Ile Met Asp Gly Gln Leu Leu Ser Gln Glu Ala Leu Glu	
645 650 655	
aaa gta gat tat cca gga aga cgt gtt ctc aca gag ctg aat agc ctc	2017
Lys Val Asp Tyr Pro Gly Arg Arg Val Leu Thr Glu Leu Asn Ser Leu	
660 665 670	
att tcc cgc ctg gcg gat gac acg aag aca tat aaa gct gag aag gct	2065
Ile Ser Arg Leu Ala Asp Asp Thr Lys Thr Tyr Lys Ala Glu Lys Ala	
675 680 685	
cgt gga gaa ttg gcg tcc agc att gaa tgt tac atg aaa gac cat cct	2113
Arg Gly Glu Leu Ala Ser Ser Ile Glu Cys Tyr Met Lys Asp His Pro	
690 695 700	
gaa tgt aca gag gaa gag gct ctc gat cac atc tat agc att ctg gag	2161
Glu Cys Thr Glu Glu Glu Ala Leu Asp His Ile Tyr Ser Ile Leu Glu	
705 710 715 720	
ccg gcg gtg aag gaa ctg aca aga gag ttt ctg aag ccc gac gac gtc	2209
Pro Ala Val Lys Glu Leu Thr Arg Glu Phe Leu Lys Pro Asp Asp Val	
725 730 735	
cca ttc gcc tgc aag aag atg ctt ttc gag gag aca aga gtg acg atg	2257
Pro Phe Ala Cys Lys Lys Met Leu Phe Glu Glu Thr Arg Val Thr Met	
740 745 750	

gtg ata ttc aag gat gga gat gga ttc ggt gtt tcc aaa tta gaa gtc 2305
Val Ile Phe Lys Asp Gly Asp Gly Phe Gly Val Ser Lys Leu Glu Val
755 760 765

aaa gat cat atc aaa gag tgt ctc att gaa ccg ctg cca ctg taa 2350
Lys Asp His Ile Lys Glu Cys Leu Ile Glu Pro Leu Pro Leu
770 775 780

tcaaaatagt tgcaataata attgaaataa tgtcaactat gtttcacaaa aaaaaaaaaa 2410
aaaaaaaaaa aaaa
2424

```
<210> 16
<211> 782
<212> PRT
<213> Abies grandis
```

<400> 16
Gly Tyr Asp Leu Val His Ser Leu Lys Ser Pro Tyr Ile Asp Ser Ser
1 5 10 15

Tyr Arg Glu Arg Ala Glu Val Leu Val Ser Glu Ile Lys Val Met Leu
20 25 30

Asn Pro Ala Ile Thr Gly Asp Gly Glu Ser Met Ile Thr Pro Ser Ala
35 40 45

Tyr Asp Thr Ala Trp Val Ala Arg Val Pro Ala Ile Asp Gly Ser Ala
50 55 60

Arg Pro Gln Phe Pro Gln Thr Val Asp Trp Ile Leu Lys Asn Gln Leu
65 70 75 80

Lys Asp Gly Ser Trp Gly Ile Gln Ser His Phe Leu Leu Ser Asp Arg
85 90 95

Leu Leu Ala Thr Leu Ser Cys Val Leu Val Leu Leu Lys Trp Asn Val
100 105 110

Gly Asp Leu Gln Val Glu Gln Gly Ile Glu Phe Ile Lys Ser Asn Leu
115 120 125

Glu Leu Val Lys Asp Glu Thr Asp Gln Asp Ser Leu Val Thr Asp Phe
130 135 140

Glu Ile Ile Phe Pro Ser Leu Leu Arg Glu Ala Gln Ser Leu Arg Leu
145 150 155 160

Gly Leu Pro Tyr Asp Leu Pro Tyr Ile His Leu Leu Gln Thr Lys Arg
165 170 175

Gln Glu Arg Leu Ala Lys Leu Ser Arg Glu Glu Ile Tyr Ala Val Pro
180 185 190

Ser Pro Leu Leu Tyr Ser Leu Glu Gly Ile Gln Asp Ile Val Glu Trp
195 200 205

Glu Arg Ile Met Glu Val Gln Ser Gln Asp Gly Ser Phe Leu Ser Ser

210	215	220
Pro Ala Ser Thr Ala Cys Val Phe Met His Thr Gly Asp Ala Lys Cys 225 230 235 240		
Leu Glu Phe Leu Asn Ser Val Met Ile Lys Phe Gly Asn Phe Val Pro 245 250 255		
Cys Leu Tyr Pro Val Asp Leu Leu Glu Arg Leu Leu Ile Val Asp Asn 260 265 270		
Ile Val Arg Leu Gly Ile Tyr Arg His Phe Glu Lys Glu Ile Lys Glu 275 280 285		
Ala Leu Asp Tyr Val Tyr Arg His Trp Asn Glu Arg Gly Ile Gly Trp 290 295 300		
Gly Arg Leu Asn Pro Ile Ala Asp Leu Glu Thr Thr Ala Leu Gly Phe 305 310 315 320		
Arg Leu Leu Arg Leu His Arg Tyr Asn Val Ser Pro Ala Ile Phe Asp 325 330 335		
Asn Phe Lys Asp Ala Asn Gly Lys Phe Ile Cys Ser Thr Gly Gln Phe 340 345 350		
Asn Lys Asp Val Ala Ser Met Leu Asn Leu Tyr Arg Ala Ser Gln Leu 355 360 365		
Ala Phe Pro Gly Glu Asn Ile Leu Asp Glu Ala Lys Ser Phe Ala Thr 370 375 380		
Lys Tyr Leu Arg Glu Ala Leu Glu Lys Ser Glu Thr Ser Ser Ala Trp 385 390 395 400		
Asn Asn Lys Gln Asn Leu Ser Gln Glu Ile Lys Tyr Ala Leu Lys Thr 405 410 415		
Ser Trp His Ala Ser Val Pro Arg Val Glu Ala Lys Arg Tyr Cys Gln 420 425 430		
Val Tyr Arg Pro Asp Tyr Ala Arg Ile Ala Lys Cys Val Tyr Lys Leu 435 440 445		
Pro Tyr Val Asn Asn Glu Lys Phe Leu Glu Leu Gly Lys Leu Asp Phe 450 455 460		
Asn Ile Ile Gln Ser Ile His Gln Glu Glu Met Lys Asn Val Thr Ser 465 470 475 480		
Trp Phe Arg Asp Ser Gly Leu Pro Leu Phe Thr Phe Ala Arg Glu Arg 485 490 495		
Pro Leu Glu Phe Tyr Phe Leu Val Ala Ala Gly Thr Tyr Glu Pro Gln 500 505 510		
Tyr Ala Lys Cys Arg Phe Leu Phe Thr Lys Val Ala Cys Leu Gln Thr 515 520 525		
Val Leu Asp Asp Met Tyr Asp Thr Tyr Gly Thr Leu Asp Glu Leu Lys		

530	535	540
Leu Phe Thr Glu Ala Val Arg Arg Trp Asp 545	550	Leu Ser Phe Thr Glu Asn 555 560
Leu Pro Asp Tyr Met Lys Leu Cys Tyr Gln Ile Tyr Tyr Asp Ile Val 565	570	575
His Glu Val Ala Trp Glu Ala Glu Lys Glu Gln Gly Arg Glu Leu Val 580	585	590
Ser Phe Phe Arg Lys Gly Trp Glu Asp Tyr Leu Leu Gly Tyr Tyr Glu 595	600	605
Glu Ala Glu Trp Leu Ala Ala Glu Tyr Val Pro Thr Leu Asp Glu Tyr 610	615	620
Ile Lys Asn Gly Ile Thr Ser Ile Gly Gln Arg Ile Leu Leu Leu Ser 625	630	635 640
Gly Val Leu Ile Met Asp Gly Gln Leu Leu Ser Gln Glu Ala Leu Glu 645	650	655
Lys Val Asp Tyr Pro Gly Arg Arg Val Leu Thr Glu Leu Asn Ser Leu 660	665	670
Ile Ser Arg Leu Ala Asp Asp Thr Lys Thr Tyr Lys Ala Glu Lys Ala 675	680	685
Arg Gly Glu Leu Ala Ser Ser Ile Glu Cys Tyr Met Lys Asp His Pro 690	695	700
Glu Cys Thr Glu Glu Glu Ala Leu Asp His Ile Tyr Ser Ile Leu Glu 705	710	715 720
Pro Ala Val Lys Glu Leu Thr Arg Glu Phe Leu Lys Pro Asp Asp Val 725	730	735
Pro Phe Ala Cys Lys Lys Met Leu Phe Glu Glu Thr Arg Val Thr Met 740	745	750
Val Ile Phe Lys Asp Gly Asp Gly Phe Gly Val Ser Lys Leu Glu Val 755	760	765
Lys Asp His Ile Lys Glu Cys Leu Ile Glu Pro Leu Pro Leu 770	775	780

<210> 17
 <211> 1967
 <212> DNA
 <213> *Abies grandis*

<220>
 <221> CDS
 <222> (3) .. (1736)
 <223> Clone AG4.30

<400> 17
 tt tct gaa tct tcc atc cct cga cgc aca ggg aat cat cac gga aat 47

	Ser 1	Glu	Ser	Ser	Ile 5	Pro	Arg	Arg	Thr	Gly 10	Asn	His	His	Gly	Asn 15	
gtg Val	tgg Trp	gac Asp	gat Asp	gac Asp 20	ctc Leu	ata Ile	cac His	tct Ser	ctc Leu 25	aac Asn	tcg Ser	ccc Pro	tat Tyr	ggg Gly 30	gca Ala	95
cct Pro	gca Ala	tat Tyr	tat Tyr 35	gag Glu	ctc Leu	ctt Leu	caa Gln	aag Lys 40	ctt Leu	att Ile	gag Glu	gag Glu	atc Ile 45	aag Lys	cat His	143
tta Leu	ctt Leu	ttg Leu 50	act Thr	gaa Glu	atg Met	gaa Glu	atg Met 55	gat Asp	gat Asp	ggc Gly	gat Asp	cat His 60	gat Asp	tta Leu	atc Ile	191
aaa Lys	cgt Arg 65	ctt Leu	cag Gln	atc Ile	gtt Val	gac Asp 70	act Thr	ttg Leu	gaa Glu	tgc Cys	ctg Leu 75	gga Gly	atc Ile	gat Asp	aga Arg	239
cat His 80	ttt Phe	gaa Glu	cac His	gaa Glu	ata Ile 85	caa Gln	aca Thr	gct Ala	gct Ala	tta Leu 90	gat Asp	tac Tyr	gtt Val	tac Tyr	aga Arg 95	287
tgg Trp	tgg Trp	aac Asn	gaa Glu	aaa Lys 100	ggt Gly	atc Ile	ggg Gly	gag Glu	gga Gly 105	tca Ser	aga Arg	gat Asp	tcc Ser	ttc Phe 110	agc Ser	335
aaa Lys	gat Asp	ctc Leu	aac Asn 115	gct Ala	aca Thr	gct Ala	tta Leu	gga Gly 120	ttt Phe	cgc Arg	gct Ala	ctc Leu	cga Arg 125	ctg Leu	cat His	383
cga Arg	tat Tyr	aac Asn 130	gta Val	tcg Ser	tca Ser	ggt Gly	gtg Val 135	ttg Leu	aag Lys	aat Asn	ttc Phe	aag Lys 140	gat Asp	gaa Glu	aac Asn	431
ggg Gly	aag Lys 145	ttc Phe	ttc Phe	tgc Cys	aac Asn	ttt Phe 150	act Thr	ggt Gly	gaa Glu	gaa Glu	gga Gly 155	aga Arg	gga Gly	gat Asp	aaa Lys	479
caa Gln 160	gtg Val	aga Arg	agc Ser	atg Met	ttg Leu 165	tcg Ser	tta Leu	ctt Leu	cga Arg	gct Ala 170	tca Ser	gag Glu	att Ile	tcg Ser	ttt Phe 175	527
ccc Pro	gga Gly	gaa Glu	aaa Lys	gtg Val 180	atg Met	gaa Glu	gag Glu	gcc Ala	aag Lys 185	gca Ala	ttc Phe	aca Thr	aga Arg	gaa Glu 190	tat Tyr	575
cta Leu	aac Asn	caa Gln	gtt Val 195	tta Leu	gct Ala	gga Gly	cac His	ggg Gly 200	gat Asp	gtg Val	act Thr	gac Asp	gtg Val 205	gat Asp	caa Gln	623
agc Ser	ctt Leu	ttg Leu 210	gag Glu	aga Arg	ggt Gly	gaa Glu	gta Val 215	cgc Arg	att Ile	gga Gly	gtt Val	tcc Ser 220	atg Met	gct Ala	tgc Cys	671
agt Ser	gtg Val 225	ccg Pro	aga Arg	tgg Trp	gag Glu	gca Ala 230	agg Arg	agc Ser	ttt Phe	ctc Leu	gaa Glu 235	ata Ile	tat Tyr	gga Gly	cac His	719
aac	cat	tcg	tgg	ctc	aag	tcg	aat	atc	aac	caa	aaa	atg	ttg	aag	tta	767

Asn 240	His	Ser	Trp	Leu	Lys 245	Ser	Asn	Ile	Asn	Gln 250	Lys	Met	Leu	Lys	Leu 255	
gcc Ala	aaa Lys	ttg Leu	gac Asp	ttc Phe	aat Asn	att Ile	ctg Leu	caa Gln	tgc Cys	aaa Lys	cat His	cac His	aag Lys	gag Glu	ata Ile	815
cag Gln	ttt Phe	att Ile	aca Thr	agg Arg	tgg Trp	tgg Trp	aga Arg	gac Asp	tcg Ser	ggt Gly	ata Ile	tcg Ser	cag Gln	ctg Leu	aat Asn	863
ttc Phe	tat Tyr	cga Arg	aag Lys	cga Arg	cac His	gtg Val	gaa Glu	tat Tyr	tat Tyr	tct Ser	tgg Trp	gtt Val	gtt Val	atg Met	tgc Cys	911
att Ile	ttt Phe	gag Glu	cca Pro	gag Glu	ttc Phe	tct Ser	gaa Glu	agt Ser	aga Arg	att Ile	gcc Ala	ttc Phe	gcc Ala	aaa Lys	act Thr	959
gct Ala	atc Ile	cta Leu	tgt Cys	act Thr	gtt Val	cta Leu	gat Asp	gac Asp	ctc Leu	tat Tyr	gat Asp	acg Thr	cac His	gca Ala	acg Thr	1007
ttg Leu	cat His	gaa Glu	atc Ile	aaa Lys	atc Ile	atg Met	aca Thr	gag Glu	gga Gly	gtg Val	aga Arg	cga Arg	tgg Trp	gat Asp	ctt Leu	1055
tcg Ser	ttg Leu	aca Thr	gat Asp	gac Asp	ctc Leu	cca Pro	gac Asp	tac Tyr	att Ile	aaa Lys	att Ile	gca Ala	ttc Phe	cag Gln	ttc Phe	1103
ttc Phe	ttc Phe	aat Asn	aca Thr	gtg Val	aat Asn	gaa Glu	ttg Leu	ata Ile	gtt Val	gaa Glu	atc Ile	gtg Val	aaa Lys	cgg Arg	caa Gln	1151
ggg Gly	cgg Arg	gat Asp	atg Met	aca Thr	acc Thr	ata Ile	gtt Val	aaa Lys	gat Asp	tgc Cys	tgg Trp	aag Lys	cga Arg	tac Tyr	att Ile	1199
gag Glu	tct Ser	tat Tyr	ctg Leu	caa Gln	gaa Glu	gcg Ala	gaa Glu	tgg Trp	ata Ile	gca Ala	act Thr	gga Gly	cat His	att Ile	ccc Pro	1247
act Thr	ttt Phe	aac Asn	gaa Glu	tac Tyr	ata Ile	aag Lys	aac Asn	ggc Gly	atg Met	gct Ala	agc Ser	tca Ser	ggg Gly	atg Met	tgt Cys	1295
att Ile	gta Val	aat Asn	ttg Leu	aat Asn	cca Pro	ctt Leu	ctc Leu	ttg Leu	ttg Leu	ggt Gly	aaa Lys	ctt Leu	ctc Leu	ccc Pro	gac Asp	1343
aac Asn	att Ile	ctg Leu	gag Glu	caa Gln	ata Ile	cat His	tct Ser	cca Pro	tcc Ser	aag Lys	atc Ile	ctg Leu	gac Asp	ctc Leu	tta Leu	1391
gaa Glu	ttg Leu	acg Thr	ggc Gly	aga Arg	atc Ile	gcc Ala	gat Asp	gac Asp	tta Leu	aaa Lys	gat Asp	ttc Phe	gag Glu	gac Asp	gag Glu	1439
aag	gaa	cgc	ggg	gag	atg	gct	tca	tct	tta	cag	tgt	tat	atg	aaa	gaa	1487

Lys Glu Arg Gly Glu Met Ala Ser Ser Leu Gln Cys Tyr Met Lys Glu
 480 485 490 495
 aat cct gaa tct aca gtg gaa aat gct tta aat cac ata aaa ggc atc 1535
 Asn Pro Glu Ser Thr Val Glu Asn Ala Leu Asn His Ile Lys Gly Ile
 500 505 510
 ctt aat cgt tcc ctt gag gaa ttt aat tgg gag ttt atg aag cag gat 1583
 Leu Asn Arg Ser Leu Glu Glu Phe Asn Trp Glu Phe Met Lys Gln Asp
 515 520 525
 agt gtc cca atg tgt tgc aag aaa ttc act ttc aat ata ggt cga gga 1631
 Ser Val Pro Met Cys Cys Lys Lys Phe Thr Phe Asn Ile Gly Arg Gly
 530 535 540
 ctt caa ttc atc tac aaa tac aga gac ggc tta tac att tct gac aag 1679
 Leu Gln Phe Ile Tyr Lys Tyr Arg Asp Gly Leu Tyr Ile Ser Asp Lys
 545 550 555
 gaa gta aag gac cag ata ttc aaa att cta gtc cac caa gtt cca atg 1727
 Glu Val Lys Asp Gln Ile Phe Lys Ile Leu Val His Gln Val Pro Met
 560 565 570 575
 gag gaa tag tgatgggtcctt ggttgtagtt gtctattatg gtatatgtgca 1776
 Glu Glu
 ttgacatttta tgcttaaagg tgtttcttaa acgttttaggg cggaccgtta aataagttgg 1836
 caataattaa tatttagaga ctttgtggaa gtgttttaggg cataaaattg cctatggcct 1896
 atggcaagct acaaattgaa attgtttgtgt ttataatatt tttattttat ttaaaaaaaaa 1956
 aaaaaaaaa a 1967

<210> 18
 <211> 577
 <212> PRT
 <213> Abies grandis

<400> 18
 Ser Glu Ser Ser Ile Pro Arg Arg Thr Gly Asn His His Gly Asn Val
 1 5 10 15
 Trp Asp Asp Asp Leu Ile His Ser Leu Asn Ser Pro Tyr Gly Ala Pro
 20 25 30
 Ala Tyr Tyr Glu Leu Leu Gln Lys Leu Ile Glu Glu Ile Lys His Leu
 35 40 45
 Leu Leu Thr Glu Met Glu Met Asp Asp Gly Asp His Asp Leu Ile Lys
 50 55 60
 Arg Leu Gln Ile Val Asp Thr Leu Glu Cys Leu Gly Ile Asp Arg His
 65 70 75 80
 Phe Glu His Glu Ile Gln Thr Ala Ala Leu Asp Tyr Val Tyr Arg Trp
 85 90 95
 Trp Asn Glu Lys Gly Ile Gly Glu Gly Ser Arg Asp Ser Phe Ser Lys

100					105					110					
Asp	Leu	Asn	Ala	Thr	Ala	Leu	Gly	Phe	Arg	Ala	Leu	Arg	Leu	His	Arg
		115					120					125			
Tyr	Asn	Val	Ser	Ser	Gly	Val	Leu	Lys	Asn	Phe	Lys	Asp	Glu	Asn	Gly
	130					135					140				
Lys	Phe	Phe	Cys	Asn	Phe	Thr	Gly	Glu	Glu	Gly	Arg	Gly	Asp	Lys	Gln
145					150					155					160
Val	Arg	Ser	Met	Leu	Ser	Leu	Leu	Arg	Ala	Ser	Glu	Ile	Ser	Phe	Pro
				165					170					175	
Gly	Glu	Lys	Val	Met	Glu	Glu	Ala	Lys	Ala	Phe	Thr	Arg	Glu	Tyr	Leu
			180					185					190		
Asn	Gln	Val	Leu	Ala	Gly	His	Gly	Asp	Val	Thr	Asp	Val	Asp	Gln	Ser
		195					200					205			
Leu	Leu	Glu	Arg	Gly	Glu	Val	Arg	Ile	Gly	Val	Ser	Met	Ala	Cys	Ser
	210					215					220				
Val	Pro	Arg	Trp	Glu	Ala	Arg	Ser	Phe	Leu	Glu	Ile	Tyr	Gly	His	Asn
225					230					235					240
His	Ser	Trp	Leu	Lys	Ser	Asn	Ile	Asn	Gln	Lys	Met	Leu	Lys	Leu	Ala
				245					250					255	
Lys	Leu	Asp	Phe	Asn	Ile	Leu	Gln	Cys	Lys	His	His	Lys	Glu	Ile	Gln
			260					265					270		
Phe	Ile	Thr	Arg	Trp	Trp	Arg	Asp	Ser	Gly	Ile	Ser	Gln	Leu	Asn	Phe
	275						280					285			
Tyr	Arg	Lys	Arg	His	Val	Glu	Tyr	Tyr	Ser	Trp	Val	Val	Met	Cys	Ile
	290					295					300				
Phe	Glu	Pro	Glu	Phe	Ser	Glu	Ser	Arg	Ile	Ala	Phe	Ala	Lys	Thr	Ala
305					310					315					320
Ile	Leu	Cys	Thr	Val	Leu	Asp	Asp	Leu	Tyr	Asp	Thr	His	Ala	Thr	Leu
				325					330					335	
His	Glu	Ile	Lys	Ile	Met	Thr	Glu	Gly	Val	Arg	Arg	Trp	Asp	Leu	Ser
			340					345					350		
Leu	Thr	Asp	Asp	Leu	Pro	Asp	Tyr	Ile	Lys	Ile	Ala	Phe	Gln	Phe	Phe
		355					360					365			
Phe	Asn	Thr	Val	Asn	Glu	Leu	Ile	Val	Glu	Ile	Val	Lys	Arg	Gln	Gly
	370					375					380				
Arg	Asp	Met	Thr	Thr	Ile	Val	Lys	Asp	Cys	Trp	Lys	Arg	Tyr	Ile	Glu
385					390					395					400
Ser	Tyr	Leu	Gln	Glu	Ala	Glu	Trp	Ile	Ala	Thr	Gly	His	Ile	Pro	Thr
				405					410					415	
Phe	Asn	Glu	Tyr	Ile	Lys	Asn	Gly	Met	Ala	Ser	Ser	Gly	Met	Cys	Ile

420 425 430
 Val Asn Leu Asn Pro Leu Leu Leu Leu Gly Lys Leu Leu Pro Asp Asn
 435 440 445
 Ile Leu Glu Gln Ile His Ser Pro Ser Lys Ile Leu Asp Leu Leu Glu
 450 455 460
 Leu Thr Gly Arg Ile Ala Asp Asp Leu Lys Asp Phe Glu Asp Glu Lys
 465 470 475 480
 Glu Arg Gly Glu Met Ala Ser Ser Leu Gln Cys Tyr Met Lys Glu Asn
 485 490 495
 Pro Glu Ser Thr Val Glu Asn Ala Leu Asn His Ile Lys Gly Ile Leu
 500 505 510
 Asn Arg Ser Leu Glu Glu Phe Asn Trp Glu Phe Met Lys Gln Asp Ser
 515 520 525
 Val Pro Met Cys Cys Lys Lys Phe Thr Phe Asn Ile Gly Arg Gly Leu
 530 535 540
 Gln Phe Ile Tyr Lys Tyr Arg Asp Gly Leu Tyr Ile Ser Asp Lys Glu
 545 550 555 560
 Val Lys Asp Gln Ile Phe Lys Ile Leu Val His Gln Val Pro Met Glu
 565 570 575

Glu

<210> 19
 <211> 1416
 <212> DNA
 <213> *Abies grandis*

<220>
 <221> CDS
 <222> (3)..(1199)
 <223> Clone AG5.9

<400> 19
 aa aaa gtg atg gaa gag gcg aag gca ttc aca aca aat tat cta aag 47
 Lys Val Met Glu Glu Ala Lys Ala Phe Thr Thr Asn Tyr Leu Lys
 1 5 10 15
 aaa gtt tta gca gga cgg gag gct acc cac gtc gat gaa agc ctt ttg 95
 Lys Val Leu Ala Gly Arg Glu Ala Thr His Val Asp Glu Ser Leu Leu
 20 25 30
 gga gag gtg aag tac gca ttg gag ttt cca tgg cat tgc agt gtg cag 143
 Gly Glu Val Lys Tyr Ala Leu Glu Phe Pro Trp His Cys Ser Val Gln
 35 40 45
 aga tgg gag gca agg agc ttt atc gaa ata ttt gga caa att gat tca 191
 Arg Trp Glu Ala Arg Ser Phe Ile Glu Ile Phe Gly Gln Ile Asp Ser
 50 55 60
 gag ctt aag tcg aat ttg agc aaa aaa atg tta gag ttg gcg aaa ttg 239

Glu 65	Leu	Lys	Ser	Asn	Leu	Ser 70	Lys	Lys	Met	Leu	Glu 75	Leu	Ala	Lys	Leu	
gac	ttc	aat	att	ctg	caa	tgc	aca	cat	cag	aaa	gaa	ctg	cag	att	atc	287
Asp 80	Phe	Asn	Ile	Leu	Gln 85	Cys	Thr	His	Gln	Lys 90	Glu	Leu	Gln	Ile	Ile 95	
tca	agg	tgg	ttc	gca	gac	tca	agt	ata	gca	tcc	ctg	aat	ttc	tat	cgg	335
Ser	Arg	Trp	Phe	Ala 100	Asp	Ser	Ser	Ile	Ala 105	Ser	Leu	Asn	Phe	Tyr 110	Arg	
aaa	tgt	tac	gtc	gaa	ttt	tac	ttt	tgg	atg	gct	gca	gcc	atc	tcc	gag	383
Lys	Cys	Tyr	Val 115	Glu	Phe	Tyr	Phe	Trp 120	Met	Ala	Ala	Ala	Ile 125	Ser	Glu	
ccg	gag	ttt	tct	gga	agc	aga	gtt	gcc	ttc	aca	aaa	att	gct	ata	ctg	431
Pro	Glu	Phe 130	Ser	Gly	Ser	Arg	Val 135	Ala	Phe	Thr	Lys	Ile 140	Ala	Ile	Leu	
atg	aca	atg	cta	gat	gac	ctg	tac	gat	act	cac	gga	acc	ttg	gac	caa	479
Met 145	Thr	Met	Leu	Asp	Asp 150	Leu	Tyr	Asp	Thr	His	Gly 155	Thr	Leu	Asp	Gln	
ctc	aaa	atc	ttt	aca	gag	gga	gtg	aga	cga	tgg	gat	gtt	tcg	ttg	gta	527
Leu 160	Lys	Ile	Phe	Thr 165	Glu	Gly	Val	Arg	Arg	Trp 170	Asp	Val	Ser	Leu	Val 175	
gag	ggc	ctc	cca	gac	ttc	atg	aaa	att	gca	ttc	gag	ttc	tgg	tta	aag	575
Glu	Gly	Leu	Pro 180	Asp	Phe	Met	Lys	Ile	Ala 185	Phe	Glu	Phe	Trp	Leu 190	Lys	
aca	tct	aat	gaa	ttg	att	gct	gaa	gct	gtt	aaa	gcg	caa	ggg	caa	gat	623
Thr	Ser	Asn 195	Glu	Leu	Ile	Ala	Glu	Ala 200	Val	Lys	Ala	Gln 205	Gly	Gln	Asp	
atg	gcg	gcc	tac	ata	aga	aaa	aat	gca	tgg	gag	cga	tac	ctt	gaa	gct	671
Met	Ala 210	Ala	Tyr	Ile	Arg	Lys	Asn 215	Ala	Trp	Glu	Arg	Tyr 220	Leu	Glu	Ala	
tat	ctg	caa	gat	gcg	gaa	tgg	ata	gcc	act	gga	cat	gtc	ccc	acc	ttt	719
Tyr 225	Leu	Gln	Asp	Ala	Glu 230	Trp	Ile	Ala	Thr	Gly 235	His	Val	Pro	Thr	Phe	
gat	gag	tac	ttg	aat	aat	ggc	aca	cca	aac	act	ggg	atg	tgt	gta	ttg	767
Asp 240	Glu	Tyr	Leu	Asn 245	Asn	Gly	Thr	Pro	Asn 250	Thr	Gly	Met	Cys	Val 255	Leu	
aat	ttg	att	ccg	ctt	ctg	tta	atg	ggt	gaa	cat	tta	cca	atc	gac	att	815
Asn	Leu	Ile	Pro 260	Leu	Leu	Leu	Met	Gly 265	Glu	His	Leu	Pro	Ile 270	Asp	Ile	
ctg	gag	caa	ata	ttc	ttg	ccc	tcc	agg	ttc	cac	cat	ctc	att	gaa	ttg	863
Leu	Glu	Gln 275	Ile	Phe	Leu	Pro	Ser 280	Arg	Phe	His	His	Leu 285	Ile	Glu	Leu	
gct	tcc	agg	ctc	gtc	gat	gac	gcg	aga	gat	ttc	cag	gcg	gag	aag	gat	911
Ala	Ser 290	Arg	Leu	Val	Asp	Asp 295	Ala	Arg	Asp	Phe	Gln 300	Ala	Glu	Lys	Asp	
cat	ggg	gat	tta	tcg	tgt	att	gag	tgt	tat	tta	aaa	gat	cat	cct	gag	959

His Gly Asp Leu Ser Cys Ile Glu Cys Tyr Leu Lys Asp His Pro Glu
 305 310 315
 tct aca gta gaa gat gct tta aat cat gtt aat ggc ctc ctt ggc aat 1007
 Ser Thr Val Glu Asp Ala Leu Asn His Val Asn Gly Leu Leu Gly Asn
 320 325 330 335
 tgc ctt ctg gaa atg aat tgg aag ttc tta aag aag cag gac agt gtg 1055
 Cys Leu Leu Glu Met Asn Trp Lys Phe Leu Lys Lys Gln Asp Ser Val
 340 345 350
 cca ctc tcg tgt aag aag tac agc ttc cat gta ttg gca cga agc atc 1103
 Pro Leu Ser Cys Lys Lys Tyr Ser Phe His Val Leu Ala Arg Ser Ile
 355 360 365
 caa ttc atg tac aat caa ggc gat ggc ttc tcc att tcg aac aaa gtg 1151
 Gln Phe Met Tyr Asn Gln Gly Asp Gly Phe Ser Ile Ser Asn Lys Val
 370 375 380
 atc aag gat caa gtg cag aaa gtt ctt att gtc ccc gtg cct att tga 1199
 Ile Lys Asp Gln Val Gln Lys Val Leu Ile Val Pro Val Pro Ile
 385 390 395
 tagtagatac tagatagtag attagtagct attagtattt atttcatatc aatatttact 1259
 aatgctgatg atgggttaaag tccattcaga ccaatctttg gtttattgga cttaaataaa 1319
 tgaattaatt agtttggtttt aaaattgtac tatttactgt tggaaataat gttttcatta 1379
 ttgaaataac tagcacaact attttagtgt gggtgat 1416

<210> 20
 <211> 398
 <212> PRT
 <213> Abies grandis

<400> 20
 Lys Val Met Glu Glu Ala Lys Ala Phe Thr Thr Asn Tyr Leu Lys Lys
 1 5 10 15
 Val Leu Ala Gly Arg Glu Ala Thr His Val Asp Glu Ser Leu Leu Gly
 20 25 30
 Glu Val Lys Tyr Ala Leu Glu Phe Pro Trp His Cys Ser Val Gln Arg
 35 40 45
 Trp Glu Ala Arg Ser Phe Ile Glu Ile Phe Gly Gln Ile Asp Ser Glu
 50 55 60
 Leu Lys Ser Asn Leu Ser Lys Lys Met Leu Glu Leu Ala Lys Leu Asp
 65 70 75 80
 Phe Asn Ile Leu Gln Cys Thr His Gln Lys Glu Leu Gln Ile Ile Ser
 85 90 95
 Arg Trp Phe Ala Asp Ser Ser Ile Ala Ser Leu Asn Phe Tyr Arg Lys
 100 105 110
 Cys Tyr Val Glu Phe Tyr Phe Trp Met Ala Ala Ala Ile Ser Glu Pro

115					120					125					
Glu	Phe	Ser	Gly	Ser	Arg	Val	Ala	Phe	Thr	Lys	Ile	Ala	Ile	Leu	Met
130						135					140				
Thr	Met	Leu	Asp	Asp	Leu	Tyr	Asp	Thr	His	Gly	Thr	Leu	Asp	Gln	Leu
145					150					155					160
Lys	Ile	Phe	Thr	Glu	Gly	Val	Arg	Arg	Trp	Asp	Val	Ser	Leu	Val	Glu
				165					170					175	
Gly	Leu	Pro	Asp	Phe	Met	Lys	Ile	Ala	Phe	Glu	Phe	Trp	Leu	Lys	Thr
			180					185					190		
Ser	Asn	Glu	Leu	Ile	Ala	Glu	Ala	Val	Lys	Ala	Gln	Gly	Gln	Asp	Met
		195					200					205			
Ala	Ala	Tyr	Ile	Arg	Lys	Asn	Ala	Trp	Glu	Arg	Tyr	Leu	Glu	Ala	Tyr
	210					215					220				
Leu	Gln	Asp	Ala	Glu	Trp	Ile	Ala	Thr	Gly	His	Val	Pro	Thr	Phe	Asp
225					230					235					240
Glu	Tyr	Leu	Asn	Asn	Gly	Thr	Pro	Asn	Thr	Gly	Met	Cys	Val	Leu	Asn
			245						250					255	
Leu	Ile	Pro	Leu	Leu	Leu	Met	Gly	Glu	His	Leu	Pro	Ile	Asp	Ile	Leu
			260					265					270		
Glu	Gln	Ile	Phe	Leu	Pro	Ser	Arg	Phe	His	His	Leu	Ile	Glu	Leu	Ala
		275					280					285			
Ser	Arg	Leu	Val	Asp	Asp	Ala	Arg	Asp	Phe	Gln	Ala	Glu	Lys	Asp	His
	290					295					300				
Gly	Asp	Leu	Ser	Cys	Ile	Glu	Cys	Tyr	Leu	Lys	Asp	His	Pro	Glu	Ser
305					310					315					320
Thr	Val	Glu	Asp	Ala	Leu	Asn	His	Val	Asn	Gly	Leu	Leu	Gly	Asn	Cys
				325					330					335	
Leu	Leu	Glu	Met	Asn	Trp	Lys	Phe	Leu	Lys	Lys	Gln	Asp	Ser	Val	Pro
			340					345					350		
Leu	Ser	Cys	Lys	Lys	Tyr	Ser	Phe	His	Val	Leu	Ala	Arg	Ser	Ile	Gln
		355					360					365			
Phe	Met	Tyr	Asn	Gln	Gly	Asp	Gly	Phe	Ser	Ile	Ser	Asn	Lys	Val	Ile
	370					375					380				
Lys	Asp	Gln	Val	Gln	Lys	Val	Leu	Ile	Val	Pro	Val	Pro	Ile		
385					390					395					

<210> 21

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide PCR primer E wherein the letter
"n" represents an inosine residue

<220>

<221> misc_feature

<222> (1)..(23)

<223> PCR primer E wherein the letter n represents
inosine

<400> 21

ggngaramrr tnatggarga rgc

23

<210> 22

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: degenerate
oligonucleotide primer F wherein the letter "n"
represents an inosine residue

<220>

<221> misc_feature

<222> (1)..(24)

<223> PCR primer F wherein the letter n represents
inosine

<400> 22

garytn Cary tnhbnmgntg gtgg

24

<210> 23

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: degenerate
oligonucleotide PCR primer G wherein the letter
"n" represents an inosine residue

<220>

<221> misc_feature

<222> (1)..(21)

<223> PCR primer G wherein the letter n represents
inosine

<400> 23

ccarttnarn ccyytnacrt c

21

<210> 24

<211> 533

<212> DNA

<213> Abies grandis

<400> 24

ggggaaaaaa tgatggagga agctgaaatc ttctctacca aatattttaa agaagccctg 60
 caaaagattc cggctctccag tctttcgcga gagatcgggg acgttttgga atatggttgg 120
 cacacatatt tgccgcgatt ggaagcaagg aattacatcc aagtctttgg acaggacact 180
 gagaacacga agtcatatgt gaagagcaaa aaacttttag aactcgcaaa attggagttc 240
 aacatctttc aatccttact cgcataatccg cattgcaacc cattctgaca atggacatcc 300
 cctttcctga tcatatcctc aaggaagttg acttcccatc aaagcttaac gacttggcat 360
 gtgccatcct tcgattacga ggtgatacgc ggtgctacaa ggcggacagg gctcgtggag 420
 aagaagcttc ctctatatca tgttatatga aagacaatcc tggagtatca gaggaagatg 480
 ctctcgatca tatcaacgcc atgatcagtg acgaagtcaa aggcttcaat tgg 533

<210> 25

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: conserved
 amino acid motif on which the sequence of Primer D
 was based, wherein Xaa at position number 3 represents Thr or Ile, Xaa at
 position number 4 represents Ile or Tyr or Phe, Xaa at position number 6
 represents Ala or Val and Xaa at position number 8 represents Ala or Gly

<220>

<221> SITE

<222> (1)..(8)

<223> conserved amino acid motif on which sequence of
 primer D was based

<400> 25

Asp Asp Xaa Xaa Asp Xaa Tyr Xaa
 1 5

<210> 26

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: conserved
 amino acid motif on which the sequence of Primer E
 was based wherein Xaa at position 3 represents Lys or Thr, Xaa at position
 4 represents Val or Ile, Xaa at position 6 represents Glu or Asp

<220>

<221> SITE

<222> (1)..(8)

<223> conserved amino acid sequence on which the
 sequence of primer E was based

<400> 26

Gly Glu Xaa Xaa Met Xaa Glu Ala
 1 5

<210> 27
 <211> 7
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: conserved
 amino acid sequence on which the sequence of
 primer F was based wherein Xaa at position 2 represents Phe or Tyr or Asp
 Xaa at position 3 represents Ile or Leu, Xaa at position 4 represents Thr or Leu
 or Arg

<220>
 <221> SITE
 <222> (1)..(7)
 <223> conserved amino acid sequence on which the
 sequence of primer F was based

<400> 27
 Gln Xaa Xaa Xaa Arg Trp Trp
 1 5

<210> 28
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: conserved
 amino acid motif on which the sequence of primer G
 was based wherein Xaa at position 6 represents Phe or Leu

<220>
 <221> SITE
 <222> (1)..(8)
 <223> conserved amino acid sequence on which the
 sequence of primer G was based

<400> 28
 Asp Val Ile Lys Gly Xaa Asn Trp
 1 5

<210> 29
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: T3 primer
 oligonucleotide sequence

<400> 29
 aattaaccct cactaaaggg

<210> 30
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: T7
 oligonucleotide primer sequence

<400> 30
 gtaatacgac tcactatagg gc

22

<210> 31
 <211> 2205
 <212> DNA
 <213> *Abies grandis*

<220>
 <221> CDS
 <222> (57)..(1943)
 <223> Clone AG3.48

<400> 31
 gttatcttga gcttcctcca tataggccaa cacatatcat atcaaaggga gcaaga atg 59
 Met
 1

gct ctg gtt tct atc tca ccg ttg gct tcg aaa tct tgc ctg cgc aag 107
 Ala Leu Val Ser Ile Ser Pro Leu Ala Ser Lys Ser Cys Leu Arg Lys
 5 10 15

tcg ttg atc agt tca att cat gaa cat aag cct ccc tat aga aca atc 155
 Ser Leu Ile Ser Ser Ile His Glu His Lys Pro Pro Tyr Arg Thr Ile
 20 25 30

cca aat ctt gga atg cgt agg cga ggg aaa tct gtc acg cct tcc atg 203
 Pro Asn Leu Gly Met Arg Arg Arg Gly Lys Ser Val Thr Pro Ser Met
 35 40 45

agc atc agt ttg gcc acc gct gca cct gat gat ggt gta caa aga cgc 251
 Ser Ile Ser Leu Ala Thr Ala Ala Pro Asp Asp Gly Val Gln Arg Arg
 50 55 60 65

ata ggt gac tac cat tcc aat atc tgg gac gat gat ttc ata cag tct 299
 Ile Gly Asp Tyr His Ser Asn Ile Trp Asp Asp Phe Ile Gln Ser
 70 75 80

cta tca acg cat tat ggg gaa ccc tct tac cag gaa cgt gct gag aga 347
 Leu Ser Thr His Tyr Gly Glu Pro Ser Tyr Gln Glu Arg Ala Glu Arg
 85 90 95

tta att gtg gag gta aag aag ata ttc aat tca atg tac ctg gat gat 395
 Leu Ile Val Glu Val Lys Lys Ile Phe Asn Ser Met Tyr Leu Asp Asp
 100 105 110

gga aga tta atg agt tcc ttt aat gat ctc atg caa cgc ctt tgg ata 443
 Gly Arg Leu Met Ser Ser Phe Asn Asp Leu Met Gln Arg Leu Trp Ile
 115 120 125

gtc Val 130	gat Asp	agc Ser	gtt Val	gaa Glu	cgt Arg 135	ttg Leu	ggg Gly	ata Ile	gct Ala	aga Arg 140	cat His	ttc Phe	aag Lys	aac Asn	gag Glu 145	491
ata Ile	aca Thr	tca Ser	gct Ala	ctg Leu 150	gat Asp	tat Tyr	gtt Val	ttc Phe 155	cgt Arg	tac Tyr	tgg Trp	gag Glu	gaa Glu	aac Asn 160	ggc Gly	539
att Ile	gga Gly	tgt Cys	ggg Gly 165	aga Arg	gac Asp	agt Ser	att Ile	gtt Val 170	act Thr	gat Asp	ctc Leu	aac Asn	tca Ser 175	act Thr	gcg Ala	587
ttg Leu	ggg Gly	ttt Phe 180	cga Arg	act Thr	ctt Leu	cga Arg	tta Leu 185	cac His	ggg Gly	tac Tyr	act Thr	gta Val 190	tct Ser	cca Pro	gag Glu	635
gtt Val 195	tta Leu	aaa Lys	gct Ala	ttt Phe	caa Gln	gat Asp 200	caa Gln	aat Asn	gga Gly	cag Gln	ttt Phe 205	gta Val	tgc Cys	tcc Ser	ccc Pro	683
ggg Gly 210	cag Gln	aca Thr	gag Glu	ggg Gly	gag Glu 215	atc Ile	aga Arg	agc Ser	gtt Val	ctt Leu 220	aac Asn	tta Leu	tat Tyr	cgg Arg	gct Ala 225	731
tcc Ser	ctc Leu	att Ile	gcc Ala	ttc Phe 230	cct Pro	ggg Gly	gag Glu	aaa Lys	gtt Val 235	atg Met	gaa Glu	gaa Glu	gct Ala	gaa Glu	atc Ile 240	779
ttc Phe	tcc Ser	aca Thr	aga Arg 245	tat Tyr	ttg Leu	aaa Lys	gaa Glu	gct Ala 250	cta Leu	caa Gln	aag Lys	att Ile	cca Pro 255	gtc Val	tcc Ser	827
gct Ala	ctt Leu	tca Ser 260	caa Gln	gag Glu	ata Ile	aag Lys	ttt Phe 265	gtt Val	atg Met	gaa Glu	tat Tyr	ggc Gly 270	tgg Trp	cac His	aca Thr	875
aat Asn 275	ttg Leu	cca Pro	aga Arg	ttg Leu	gaa Glu	gca Ala 280	aga Arg	aat Asn	tac Tyr	ata Ile	gac Asp 285	aca Thr	ctt Leu	gag Glu	aaa Lys	923
gac Asp 290	acc Thr	agt Ser	gca Ala	tgg Trp	ctc Leu 295	aat Asn	aaa Lys	aat Asn	gct Ala	ggg Gly 300	aag Lys	aag Lys	ctt Leu	tta Leu	gaa Glu 305	971
ctt Leu	gca Ala	aaa Lys	ttg Leu 310	gag Glu	ttc Phe	aat Asn	ata Ile	ttt Phe 315	aac Asn	tcc Ser	tta Leu	caa Gln	caa Gln	aag Lys 320	gaa Glu	1019
tta Leu	caa Gln	tat Tyr 325	ctt Leu	ttg Leu	aga Arg	tgg Trp	tgg Trp	aaa Lys 330	gag Glu	tcg Ser	gat Asp	ttg Leu	cct Pro 335	aaa Lys	ttg Leu	1067
aca Thr	ttt Phe 340	gct Ala	cgg Arg	cat His	cgt Arg	cat His	gtg Val 345	gaa Glu	ttc Phe	tac Tyr	act Thr	ttg Leu 350	gcc Ala	tct Ser	tgt Cys	1115
att Ile 355	gcc Ala	att Ile	gac Asp	cca Pro	aaa Lys	cat His 360	tct Ser	gca Ala	ttc Phe	aga Arg	cta Leu 365	ggc Gly	ttc Phe	gcc Ala	aaa Lys	1163

atg tgt cat ctt gtc aca gtt ttg gac gat att tac gac act ttt gga	1211
Met Cys His Leu Val Thr Val Leu Asp Asp Ile Tyr Asp Thr Phe Gly	
370 375 380 385	
acg att gac gag ctt gaa ctc ttc aca tct gca att aag aga tgg aat	1259
Thr Ile Asp Glu Leu Glu Leu Phe Thr Ser Ala Ile Lys Arg Trp Asn	
390 395 400	
tca tca gag ata gaa cac ctt cca gaa tat atg aaa tgt gtg tac atg	1307
Ser Ser Glu Ile Glu His Leu Pro Glu Tyr Met Lys Cys Val Tyr Met	
405 410 415	
gtc gtg ttt gaa act gta aat gaa ctg aca cga gag gcg gag aag act	1355
Val Val Phe Glu Thr Val Asn Glu Leu Thr Arg Glu Ala Glu Lys Thr	
420 425 430	
caa ggg aga aac act ctc aac tat gtt cga aag gct tgg gag gct tat	1403
Gln Gly Arg Asn Thr Leu Asn Tyr Val Arg Lys Ala Trp Glu Ala Tyr	
435 440 445	
ttt gat tca tat atg gaa gaa gca aaa tgg atc tct aat ggt tat ctg	1451
Phe Asp Ser Tyr Met Glu Glu Ala Lys Trp Ile Ser Asn Gly Tyr Leu	
450 455 460 465	
cca acg ttt gaa gag tac cat gag aat ggg aaa gtg agc tct gca tat	1499
Pro Thr Phe Glu Thr Tyr His Glu Asn Gly Lys Val Ser Ser Ala Tyr	
470 475 480	
cgc gta gca aca ttg caa ccc atc ctc act ttg aat gca tgg ctt cct	1547
Arg Val Ala Thr Leu Gln Pro Ile Leu Thr Leu Asn Ala Trp Leu Pro	
485 490 495	
gat tac atc ttg aag gga att gat ttt cca tcc agg ttc aat gat ttg	1595
Asp Tyr Ile Leu Lys Gly Ile Asp Phe Pro Ser Arg Phe Asn Asp Leu	
500 505 510	
gca tcg tcc ttc ctt cgg cta cga ggt gac aca cgc tgc tac aag gcc	1643
Ala Ser Ser Phe Leu Arg Leu Arg Gly Asp Thr Arg Cys Tyr Lys Ala	
515 520 525	
gat agg gat cgt ggt gaa gaa gct tcg tgt ata tca tgt tat atg aaa	1691
Asp Arg Asp Arg Gly Glu Glu Ala Ser Cys Ile Ser Cys Tyr Met Lys	
530 535 540 545	
gac aat cct gga tca acc gaa gaa gat gcc ctc aat cat atc aat gcc	1739
Asp Asn Pro Gly Ser Thr Glu Glu Asp Ala Leu Asn His Ile Asn Ala	
550 555 560	
atg gtc aat gac ata atc aaa gaa tta aat tgg gaa ctt cta aga tcc	1787
Met Val Asn Asp Ile Ile Lys Glu Leu Asn Trp Glu Leu Leu Arg Ser	
565 570 575	
aac gac aat att cca atg ctg gcc aag aaa cat gct ttt gac ata aca	1835
Asn Asp Asn Ile Pro Met Leu Ala Lys Lys His Ala Phe Asp Ile Thr	
580 585 590	
aga gct ctc cac cat ctc tac ata tat cga gat ggc ttt agt gtt gcc	1883
Arg Ala Leu His His Leu Tyr Ile Tyr Arg Asp Gly Phe Ser Val Ala	
595 600 605	

aac aag gaa aca aaa aaa ttg gtt atg gaa aca ctc ctt gaa tct atg 1931
 Asn Lys Glu Thr Lys Lys Leu Val Met Glu Thr Leu Leu Glu Ser Met
 610 615 620 625
 ctt ttt taa cta taaccatata cataataata agctcataat gctaaattat 1983
 Leu Phe
 tggccttatg acatagttta tgtatgtact tgtgtgaatt caatcatata gtgtgggtat 2043
 gattaataaag ctagagctta ctaggttagt aacatggtga taaaagttat aaaatgtgag 2103
 ttatagagat acccatgttg aataatgaat tacaaaaaga gaaatttatg tagaataaga 2163
 ttggaagctt ttcaattggt ttaaaaaaaaa aaaaaaaaaa aa 2205

<210> 32
 <211> 627
 <212> PRT
 <213> Abies grandis

<400> 32
 Met Ala Leu Val Ser Ile Ser Pro Leu Ala Ser Lys Ser Cys Leu Arg
 1 5 10 15
 Lys Ser Leu Ile Ser Ser Ile His Glu His Lys Pro Pro Tyr Arg Thr
 20 25 30
 Ile Pro Asn Leu Gly Met Arg Arg Arg Gly Lys Ser Val Thr Pro Ser
 35 40 45
 Met Ser Ile Ser Leu Ala Thr Ala Ala Pro Asp Asp Gly Val Gln Arg
 50 55 60
 Arg Ile Gly Asp Tyr His Ser Asn Ile Trp Asp Asp Asp Phe Ile Gln
 65 70 75 80
 Ser Leu Ser Thr His Tyr Gly Glu Pro Ser Tyr Gln Glu Arg Ala Glu
 85 90 95
 Arg Leu Ile Val Glu Val Lys Lys Ile Phe Asn Ser Met Tyr Leu Asp
 100 105 110
 Asp Gly Arg Leu Met Ser Ser Phe Asn Asp Leu Met Gln Arg Leu Trp
 115 120 125
 Ile Val Asp Ser Val Glu Arg Leu Gly Ile Ala Arg His Phe Lys Asn
 130 135 140
 Glu Ile Thr Ser Ala Leu Asp Tyr Val Phe Arg Tyr Trp Glu Glu Asn
 145 150 155 160
 Gly Ile Gly Cys Gly Arg Asp Ser Ile Val Thr Asp Leu Asn Ser Thr
 165 170 175
 Ala Leu Gly Phe Arg Thr Leu Arg Leu His Gly Tyr Thr Val Ser Pro
 180 185 190
 Glu Val Leu Lys Ala Phe Gln Asp Gln Asn Gly Gln Phe Val Cys Ser

195					200					205					
Pro	Gly	Gln	Thr	Glu	Gly	Glu	Ile	Arg	Ser	Val	Leu	Asn	Leu	Tyr	Arg
	210					215					220				
Ala	Ser	Leu	Ile	Ala	Phe	Pro	Gly	Glu	Lys	Val	Met	Glu	Glu	Ala	Glu
225					230					235					240
Ile	Phe	Ser	Thr	Arg	Tyr	Leu	Lys	Glu	Ala	Leu	Gln	Lys	Ile	Pro	Val
				245					250					255	
Ser	Ala	Leu	Ser	Gln	Glu	Ile	Lys	Phe	Val	Met	Glu	Tyr	Gly	Trp	His
			260					265					270		
Thr	Asn	Leu	Pro	Arg	Leu	Glu	Ala	Arg	Asn	Tyr	Ile	Asp	Thr	Leu	Glu
		275					280					285			
Lys	Asp	Thr	Ser	Ala	Trp	Leu	Asn	Lys	Asn	Ala	Gly	Lys	Lys	Leu	Leu
	290					295					300				
Glu	Leu	Ala	Lys	Leu	Glu	Phe	Asn	Ile	Phe	Asn	Ser	Leu	Gln	Gln	Lys
305					310					315					320
Glu	Leu	Gln	Tyr	Leu	Leu	Arg	Trp	Trp	Lys	Glu	Ser	Asp	Leu	Pro	Lys
				325					330					335	
Leu	Thr	Phe	Ala	Arg	His	Arg	His	Val	Glu	Phe	Tyr	Thr	Leu	Ala	Ser
			340					345					350		
Cys	Ile	Ala	Ile	Asp	Pro	Lys	His	Ser	Ala	Phe	Arg	Leu	Gly	Phe	Ala
		355					360					365			
Lys	Met	Cys	His	Leu	Val	Thr	Val	Leu	Asp	Asp	Ile	Tyr	Asp	Thr	Phe
	370					375					380				
Gly	Thr	Ile	Asp	Glu	Leu	Glu	Leu	Phe	Thr	Ser	Ala	Ile	Lys	Arg	Trp
385					390					395					400
Asn	Ser	Ser	Glu	Ile	Glu	His	Leu	Pro	Glu	Tyr	Met	Lys	Cys	Val	Tyr
			405						410					415	
Met	Val	Val	Phe	Glu	Thr	Val	Asn	Glu	Leu	Thr	Arg	Glu	Ala	Glu	Lys
			420					425					430		
Thr	Gln	Gly	Arg	Asn	Thr	Leu	Asn	Tyr	Val	Arg	Lys	Ala	Trp	Glu	Ala
		435					440					445			
Tyr	Phe	Asp	Ser	Tyr	Met	Glu	Glu	Ala	Lys	Trp	Ile	Ser	Asn	Gly	Tyr
	450					455					460				
Leu	Pro	Thr	Phe	Glu	Glu	Tyr	His	Glu	Asn	Gly	Lys	Val	Ser	Ser	Ala
465					470					475					480
Tyr	Arg	Val	Ala	Thr	Leu	Gln	Pro	Ile	Leu	Thr	Leu	Asn	Ala	Trp	Leu
				485					490					495	
Pro	Asp	Tyr	Ile	Leu	Lys	Gly	Ile	Asp	Phe	Pro	Ser	Arg	Phe	Asn	Asp
			500					505					510		
Leu	Ala	Ser	Ser	Phe	Leu	Arg	Leu	Arg	Gly	Asp	Thr	Arg	Cys	Tyr	Lys

515 520 525
 Ala Asp Arg Asp Arg Gly Glu Glu Ala Ser Cys Ile Ser Cys Tyr Met
 530 535 540
 Lys Asp Asn Pro Gly Ser Thr Glu Glu Asp Ala Leu Asn His Ile Asn
 545 550 555 560
 Ala Met Val Asn Asp Ile Ile Lys Glu Leu Asn Trp Glu Leu Leu Arg
 565 570 575
 Ser Asn Asp Asn Ile Pro Met Leu Ala Lys Lys His Ala Phe Asp Ile
 580 585 590
 Thr Arg Ala Leu His His Leu Tyr Ile Tyr Arg Asp Gly Phe Ser Val
 595 600 605
 Ala Asn Lys Glu Thr Lys Lys Leu Val Met Glu Thr Leu Leu Glu Ser
 610 615 620
 Met Leu Phe
 625

<210> 33
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR
 oligonucleotide primer 2.2 BamHI

<400> 33
 caaagggatc cagaatggct ctgg

24

<210> 34
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR
 oligonucleotide primer 2.2 Not I

<400> 34
 agtaagcggc cgctttttaa tcataccac

30

<210> 35
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR
 oligonucleotide primer 3.18 EcoRI

<400> 35

ctgcaggaat tcggcacgag c 21

<210> 36
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR
oligonucleotide primer 3.18 SmaI

<400> 36
catagccccg ggcatagatt tgagctg 27

<210> 37
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR
oligonucleotide primer 10 NdeI

<400> 37
ggcaggaaca tatggctctc ctttctatcg 30

<210> 38
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR
oligonucleotide primer 10 BamHI

<400> 38
tctagaacta gtggatcccc cgggctgcag 30

<210> 39
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR
oligonucleotide primer JB29

<400> 39
ctaccattcc aatatctg 18

<210> 40
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR
oligonucleotide primer 2-8

<400> 40
gttggatctt agaagttccc 20

<210> 41
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR
oligonucleotide primer 3-9

<400> 41
tttccattcc aacctctggg 20

<210> 42
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR
oligonucleotide primer 3-11

<400> 42
cgtaatggaa agctctggcg 20

<210> 43
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR
oligonucleotide primer 7-1

<400> 43
ccttacacgc ctttgatgg 20

<210> 44
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR
oligonucleotide sequence 7-3

<400> 44
tctgttgatc caggatggtc 20

<210> 45
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: conserved
amino acid motif common to all prenyl transferases wherein Xaa at position
3 and 4 represents any amino acid

<400> 45
Asp Asp Xaa Xaa Asp
1 5

<210> 46
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: amino acid
motif from which oligonucleotide primers can be
synthesized that hybridize to the monoterpene
synthases of the present invention, wherein Xaa at position 4 represents
Leu or Ile or Val

<400> 46
His Ser Asn Xaa Trp Asp Asp Asp
1 5

<210> 47
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: amino acid
motif from which degenerate oligonucleotides can
be constructed that hybridize to the monoterpene
synthases of the present invention

<400> 47
Ala Leu Asp Tyr Val Tyr
1 5

<210> 48
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: amino acid
motif from which degenerate oligonucleotide
sequences can be constructed that hybridize to the
monoterpene synthases of the present invention

<400> 48

Glu Leu Ala Lys Leu Glu Phe
1 5

<210> 49
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: amino acid
motif from which degenerate oligonucleotide
sequences can be constructed that hybridize to
monoterpene synthase clones of the present
invention

<400> 49
Arg Trp Trp Lys Glu Ser
1 5

<210> 50
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: amino acid
motif from which oligonucleotide sequences can be
constructed that hybridize to monoterpene synthase
clones of the present invention, wherein Xaa at position 1 represents Val
or Ile or Leu

<400> 50
Xaa Leu Asp Asp Met Tyr Asp
1 5

<210> 51
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: amino acid
motif from which oligonucleotide sequences can be
constructed that hybridize to monoterpene synthase
clones of the present invention wherein Xaa at position 1 represents Val
or Ile or Leu

<400> 51
Xaa Leu Asp Asp Leu Tyr Asp
1 5

<210> 52
<211> 7
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: amino acid motif from which oligonucleotide sequences can be constructed that hybridize to the monoterpene synthase clones of the present invention, wherein Xaa at position 1 represents Val or Ile or Leu

<400> 52

Xaa Leu Asp Asp Ile Tyr Asp
1 5

<210> 53

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: amino acid motif from which oligonucleotide sequences can be constructed that hybridize to the monoterpene synthase clones of the present invention, wherein Xaa at position 6 represents Asn or His

<400> 53

Cys Tyr Met Lys Asp Xaa Pro
1 5

<210> 54

<211> 9

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: exemplary oligonucleotide that corresponds to peptide sequence MetMetMet

<400> 54

atgatgatg

9

<210> 55

<211> 9

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: exemplary oligonucleotide sequence that corresponds to peptide sequence MetMetMet

<400> 55

tactactac

9

<210> 56

<211> 9

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: exemplary
oligonucleotide that corresponds to peptide
sequence MetMetMet, n is inosine

<400> 56

nacnacnac

9

<210> 57

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide corresponding to amino acid
sequence set forth in SEQ ID NO:46

<220>

<221> misc_feature

<222> (1)..(24)

<223> Oligonucleotide that corresponds to the conserved
amino acid sequence set forth in SEQ ID NO:46

<400> 57

gtgtcgttgg agaccctgct gctg

24

<210> 58

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide sequence corresponding to amino
acid sequence set forth in SEQ ID NO:47

<220>

<221> misc_feature

<222> (1)..(18)

<223> Oligonucleotide corresponding to amino acid
sequence set forth in SEQ ID NO:47

<400> 58

cgggagctga tgcagatg

18

<210> 59

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide that corresponds to amino acid
sequence set forth in SEQ ID NO:48

<220>
<221> misc_feature
<222> (1)..(21)
<223> Oligonucleotide that corresponds to conserved
amino acid sequence set forth in SEQ ID NO:48

<400> 59
ctcgagcggg tcgagctcaa g

21

<210> 60
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide that corresponds to amino acid
sequence set forth in SEQ ID NO:49

<220>
<221> misc_feature
<222> (1)..(18)
<223> Oligonucleotide that corresponds to conserved
amino acid sequence set forth in SEQ ID NO:49

<400> 60
gccaccacct tcctctcg

18

<210> 61
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide sequence corresponding to amino
acid sequence set forth in SEQ ID NO:50

<220>
<221> misc_feature
<222> (1)..(21)
<223> Oligonucleotide sequence corresponding to amino
acid sequence set forth in SEQ ID NO:50

<400> 61
gaggagctgc tgtacatgct g

21

<210> 62
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide corresponding to amino acid
sequence set forth in SEQ ID NO:51

<220>
<221> misc_feature
<222> (1)..(21)
<223> Oligonucleotide corresponding to conserved amino
acid sequence set forth in SEQ ID NO:51

<400> 62
gaggagctgc tggagatgct g

21

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/14528

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 426/601, 622; 435/320.1, 419; 530/370; 536/23.6; 800/281, 298

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 426/601, 622; 435/320.1, 419; 530/370; 536/23.6; 800/281, 298

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 95/11913 A1 (WASHINGTON STATE UNIVERSITY RESEARCH FOUNDATION) 04 May 1995, see page 3, lines 13-35 and pages 32-41.	1,3,5,7,10,13, 15-22,24,26-46, 49,50
Y	WILDUNG et al. A cDNA Clone for Taxadiene Synthase, the Diterpene Cyclase That Catalyzes the Committed Step of Taxol Biosynthesis. The Journal of Biological Chemistry. 19 April 1996. Vol. 271, No. 16, pages 9201-9204, see especially pages 9201-9202.	1,3,5,7,10, 13, 15-22,24,26-46, 49,50

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

26 OCTOBER 1998

Date of mailing of the international search report

10 NOV 1998

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

THANDA WAI

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/14528

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	COLBY et al. 4S-Limonene Synthase from the Oil Glands of Spearmint (<i>Mentha spicata</i>). The Journal of Biological Chemistry. 05 November 1993. Vol. 268, No. 31, pages 23016-23024, see especially pages 23017-23018.	1,3,5,7,10, 13, 15-22,24,26-46, 49,50
Y	YUBA et al. cDNA Cloning, Characterization, and Functional Expression of 4S(-)-Limonene Synthase from <i>Perilla frutescens</i> . Archives of Biochemistry and Biophysics. 15 August 1996. Vol. 332, No. 21, pages 280287, see especially pages 281-282.	1,3,5,7,10,13, 15-22,24,26-46, 49,50
A,P	BOHLMANN et al. Terpenoid-based defenses in conifers: cDNA cloning, characterization, and functional expression of wound-inducible (E)-alpha-bisabolene synthase from grand fir (<i>Abies grandis</i>). PNAS USA. June 1998. Vol 95, No. 12, pages 6756-6761, see entire document.	1-46,49,50
A, P	BOHLMANN et al. Monoterpene Synthases from Grand Fir (<i>Abies grandis</i>): cDNA Isolation, Characterization, and Functional Expression of Myrcene Synthase, (-)-(4S)-Limonene Synthase, and (-)-(1S,5S)-Pinene Synthase. The Journal of Biological Chemistry. 29 August 1997, Vol. 272, No. 35, pages 21784-21792, see entire document.	1-50
Y	US 4,948,811 A (SPINNER et al.) 14 August 1990) see column 2, lines 20-54 and column 3, lines 57-50.	47, 48

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/14528

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A01H 5/00, 5/10; A23D 7/00; A23K 1/14; C07H 21/04; C07K 4/10; C12N 15/04, 15/63, 15/82

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, AGRICOLA, BIOSIS, EMBASE, WPIDS, MPSEARCH (for SEQ ID NO:1-6 and 46-53)

search terms: monoterpene synthase#, myrcene synthase#, pinene synthase#, gymnosperm?, grand fir, *Abies grandis*, *A. grandis*, oil, meal#

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1, 2, 5, 6, 9, 12, 15-17, 26-46, 49, and 50, drawn to nucleotide sequences encoding myrcene synthase from Grand fir (*Abies grandis*), vectors comprising the nucleotide sequence, plants transformed with the vectors, seeds made by the transformed plants, and to a method of using the nucleotide sequence, the first product, the first method of making the product, and the first method of using the first product.

Group II, claims 1, 3, 5, 7, 10, 13, 15-17, 26-46, 49, and 50, drawn to nucleotide sequences encoding limonene synthase from Grand fir (*Abies grandis*), vectors comprising the nucleotide sequence, plants transformed with the vectors, and seeds made by the transformed plants, the second product.

Group III, claims 1, 4, 5, 8, 11, 14, 15-17, 26-46, 49, and 50, drawn to nucleotide sequences encoding pinene synthase from Grand fir (*Abies grandis*), vectors comprising the nucleotide sequence, plants transformed with the vectors, and seeds made by the transformed plants, the third product.

Group IV, claims 18-22 and 23, drawn to an isolated myrcene synthase protein from Grand fir (*Abies grandis*), the fourth product.

Group V, claims 18-22 and 24, drawn to an isolated limonene synthase protein from Grand fir (*Abies grandis*), the fifth product.

Group VI, claims 18-22 and 25, drawn to an isolated myrcene synthase protein from Grand fir (*Abies grandis*), the sixth product.

Group VII, claims 15-17, 26-46, 49, and 50, drawn to a method of using nucleotide sequences encoding limonene synthase from Grand fir (*Abies grandis*), the second method of using the second product.

Group VIII, claims 15-17, 26-46, 49, and 50, drawn to a method of using nucleotide sequences encoding pinene synthase from Grand fir (*Abies grandis*), the third method of using the third product.

Group IX, claims 47 and 48, drawn to oil and meal extracted from the seeds of Groups I-III, the seventh product.

The inventions listed as Groups I-IX do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the products of Groups I-VI and IX are distinct. The methods of Groups VII and VIII do not utilize the product of Group I. PCT Rule 13 does not provide for multiple methods of using within a single application (37 CFR 1.475(d)).

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/14528

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

